

FYJC - MATHEMATICS & STATISTICS

HIGHLIGHTS

- ✓ *Solution to all questions*
- ✓ *solutions are put in way the student is expected to reproduce in the exam*
- ✓ *taught in the class room the same way as the solution are put up here . That makes the student to easily go through the solution & prepare him/herself when he/she sits back to revise and recall the topic at any given point of time .*
- ✓ *lastly, if student due to some unavoidable reasons , has missed the lecture , will not have to run here and there to update his/her notes .*
- ✓ *however class room lectures are must for easy passage of understanding & learning the minuest details of the given topic*

PAPER - I

DETERMINANTS

Q SET - 1

CRAMER'S RULE

01. $2x - y + 3z = 9$
 $x + y + z = 6$
 $x - y + z = 2$ SS : {1,2,3}
02. $x + y + z = 6$
 $x - y + z = 2$
 $2x + y - z = 1$ SS : {1,2,3}
03. $x - y + z = 4$
 $2x + y - 3z = 0$
 $x + y + z = 2$ SS : {2,-1,1}
04. $x + y + z = 2$
 $x + 2y + z = 1$
 $5x + y + z = 6$ SS : {1,-1,2}
05. $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 2$
 $\frac{1}{x} - \frac{2}{y} + \frac{1}{z} = 3$
 $\frac{2}{x} - \frac{1}{y} + \frac{3}{z} = -1$ SS : {1,-1,2}
06. $\sin x + \cos y + \tan z = 3$
 $2\sin x + \cos y + \tan z = 4$
 $3\sin x + 4\cos y - 2\tan z = 5$
 SS : $\{\pi/2, 0, \pi/4\}$
07. $5e^x + 4\log_{10}y - 3\sqrt{z} = 1$
 $4e^x + 3\log_{10}y - 2\sqrt{z} = 2$
 $e^x + 2\log_{10}y - \sqrt{z} = 1$
 SS { 0 , 100 , 16 }
08. sum of 3 numbers is 2 . If twice the second number is added to sum of first and third we get 1 . On adding sum of second and third number to 5 times the first number we get 6 . Find the three numbers
 SS : {1,-1,2}

PROPERTIES OF DETERMINANTS

Q SET - 2

WITHOUT EXPANSION
SHOW THAT

01. $\begin{vmatrix} 1 & 1 & x \\ 1 & x & x^2 \\ 1 & x^2 & x^3 \end{vmatrix} = 0$
02. $\begin{vmatrix} x+a & x+b & x+c \\ y+a & y+b & y+c \\ z+a & z+b & z+c \end{vmatrix} = 0$
03. $\begin{vmatrix} x-y & x+y & x \\ z-x & z+x & z \\ y-z & y+z & y \end{vmatrix} = 0$
04. $\begin{vmatrix} a-b & b-c & c-a \\ b-c & c-a & a-b \\ c-a & a-b & b-c \end{vmatrix} = 0$
05. $\begin{vmatrix} 1 & a & b+c \\ 1 & b & c+a \\ 1 & c & a+b \end{vmatrix} = 0$
06. $\begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ y+z & z+x & x+y \end{vmatrix} = 0$
07. $\begin{vmatrix} 1 & xy & z(x+y) \\ 1 & yz & x(y+z) \\ 1 & zx & y(z+x) \end{vmatrix} = 0$
08. $\begin{vmatrix} xa & yb & zc \\ a^2 & b^2 & c^2 \\ 1 & 1 & 1 \end{vmatrix} = \begin{vmatrix} x & y & z \\ a & b & c \\ bc & ca & ab \end{vmatrix}$
09. $\begin{vmatrix} lp & mq & nr \\ p^2 & q^2 & r^2 \\ 1 & 1 & 1 \end{vmatrix} = \begin{vmatrix} l & m & n \\ p & q & r \\ qr & pr & pq \end{vmatrix}$
10. $\begin{vmatrix} bc & a & a^2 \\ ca & b & b^2 \\ ab & c & c^2 \end{vmatrix} = \begin{vmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{vmatrix}$

$$11. \begin{vmatrix} 1 & x^2 & x^3 \\ 1 & y^2 & y^3 \\ 1 & c^2 & c^3 \end{vmatrix} = \begin{vmatrix} yz & x & x^2 \\ xz & y & y^2 \\ xy & z & z^2 \end{vmatrix}$$

$$03. \begin{vmatrix} 1 & a & bc \\ 1 & b & ca \\ 1 & c & ab \end{vmatrix} = (a-b)(b-c)(c-a)$$

$$12. \begin{vmatrix} x+y & y+z & z+x \\ z+x & x+y & y+z \\ y+z & z+x & x+y \end{vmatrix} = 2 \begin{vmatrix} x & y & z \\ z & x & y \\ y & z & x \end{vmatrix}$$

$$04. \begin{vmatrix} a & b & c \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{vmatrix} = abc(a-b)(b-c)(c-a)$$

$$13. \begin{vmatrix} a+b & b+c & c+a \\ x+y & y+z & z+x \\ p+q & q+r & r+p \end{vmatrix} = 2 \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix}$$

$$05. \begin{vmatrix} x & p & q \\ p & x & q \\ p & q & x \end{vmatrix} = (x-p)(x-q)(x+p+q)$$

$$14. \begin{vmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{vmatrix} = 0$$

$$06. \begin{vmatrix} a & b+c & a^2 \\ b & c+a & b^2 \\ c & a+b & c^2 \end{vmatrix} = -(a-b)(b-c)(c-a)(a+b+c)$$

$$15. \begin{vmatrix} 0 & x-y & y-z \\ y-x & 0 & z-x \\ z-y & x-z & 0 \end{vmatrix} = 0$$

$$07. \begin{vmatrix} a+b+2c & a & b \\ c & b+c+2a & b \\ c & a & c+a+2b \end{vmatrix} = 2(a+b+c)^3$$

$$16. \begin{vmatrix} 11 & 4 & 10 \\ 2 & 7 & 6 \\ 5 & 1 & 4 \end{vmatrix} = 3 \begin{vmatrix} 5 & 2 & 3 \\ 11 & 5 & 2 \\ 10 & 4 & 6 \end{vmatrix}$$

$$08. \begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ x^3 & y^3 & z^3 \end{vmatrix} = (x-y)(y-z)(z-x)(x+y+z)$$

PROPERTIES OF DETERMINANTS

Q SET - 3

**WITHOUT EXPANDING
AS FAR AS POSSIBLE**

$$01. \begin{vmatrix} a^2+2a & 2a+1 & 1 \\ 2a+1 & a+2 & 1 \\ 3 & 3 & 1 \end{vmatrix} = (a-1)^3$$

10. WITHOUT EXPANSION, PROVE

$$\begin{vmatrix} 6 & 1 & 2 \\ 7 & 2 & 3 \\ 2 & 3 & -4 \end{vmatrix} + \begin{vmatrix} 2 & 4 & 7 \\ 3 & 4 & 2 \\ -4 & 3 & 3 \end{vmatrix} = 5 \begin{vmatrix} 2 & 2 & 7 \\ 3 & 1 & 2 \\ -4 & 1 & 3 \end{vmatrix}$$

$$02. \begin{vmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{vmatrix} = (x-y)(y-z)(z-x)$$

11. WITHOUT EXPANSION, PROVE

$$\begin{vmatrix} 2 & -3 & 4 \\ 5 & -6 & 2 \\ -3 & 1 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 5 & -3 \\ 5 & 8 & 5 \\ 4 & 2 & 3 \end{vmatrix} = -2 \begin{vmatrix} 2 & 4 & 1 \\ 5 & 2 & 1 \\ -3 & 3 & 3 \end{vmatrix}$$

Q SET - 4

CONSISTENCY OF EQUATIONS

01. $x + y = 3$; $5x + 6y = 17$; $2x - 3y = k$
are consistent . Find k
02. $3x + y = 2$; $kx + 2y = 3$; $2x - y = -3$
are consistent . Find k
03. $x + 3y + 2 = 0$; $4y + 2x = k$; $x - 2y = 3k$.
are consistent . Find k
04. $2x - y + 3 = 0$; $7x - 2y + 2 = 0$;
 $kx - y - 1 = 0$ are consistent . Find k & SS
05. $3x + y = 2$; $kx + 2y = 3$; $2x - y = -3$.
Find k if the lines are concurrent . Hence
find the point of concurrency
06. $kx + 3y + 4 = 0$; $x + ky + 3 = 0$;
 $3x + 4y + 5 = 0$ are consistent .
Find k . Hence find common solution set
for smallest value of k
07. if the equations $ax + by + c = 0$;
 $cx + ay + b = 0$; and $bx + cy + a = 0$
are consistent in x and y then show that
 $a^3 + b^3 + c^3 = 3abc$
08. if $l = \frac{y-1}{x}$; $m = \frac{1-x}{y}$; $n = x-y$
are consistent in x and y then prove :
 $l + m + n + lmn = 0$

QSET 5

AREA OF TRIANGLE

- Q1.** Find area of triangle with vertices as
01. (4,5) ; (0,7) ; (-1,1)
02. (3,6) ; (-1,3) ; (2,-1)
03. (1,1) , (3,7) , (10,8)
- Q2.**
01. Find k if the area of the triangle whose vertices are A (4 , k) ; B (-5 , -7); C(-4 , 1) is 38 sq. units
02. Find k if the area of the triangle whose vertices are P (k , -4) ; Q (1 , -2) ; R (4 , -5) is $15\frac{1}{2}$ sq. units
03. Find k if the area of the triangle whose vertices are P (3 , -5) ; Q (-2 , k) ; R (1 , 4) is $33\frac{1}{2}$ sq. units
- Q3.** Find area of Quadrilateral whose vertices are A (2 , 1) ; B (2 , 3) ; C(-2 , 2) ; D(-1 , 0)
- Q4.**
Using determinants show that the following set of points are collinear
01. A(3,1) ; B(4,2) ; C(5, 3)
02. A(1,-2) ; B(3,1) ; C(5,4)
03. A(3,7) ; B(4,-3) ; C(5,-13)
04.
points A(a,0) ; B(0,b) , C(1,1) are collinear .
Prove : $\frac{1}{a} + \frac{1}{b} = 1$
- 05
if points A(a ,b) ; B(c , d) ; C(a - c , b - d) are collinear then prove that $ad - bc = 0$

SOLUTION - QSET 1

01. $2x - y + 3z = 9$

$$x + y + z = 6$$

$$x - y + z = 2$$

$$D = \begin{vmatrix} + & - & + \\ 2 & -1 & 3 \\ 1 & 1 & 1 \\ 1 & -1 & 1 \end{vmatrix} = 2(1+1)+1(1-1)+3(-1-1)$$

$$= 2(2) + 1(0) + 3(-2)$$

$$= 4-6$$

$$= -2$$

$$D_x = \begin{vmatrix} + & - & + \\ 9 & -1 & 3 \\ 6 & 1 & 1 \\ 2 & -1 & 1 \end{vmatrix} = 9(1+1)+1(6-2)+3(-6-2)$$

$$= 9(2) + 1(4) + 3(-8)$$

$$= 18 + 4 - 24$$

$$= 22 - 24$$

$$= -2$$

$$D_y = \begin{vmatrix} + & - & + \\ 2 & 9 & 3 \\ 1 & 6 & 1 \\ 1 & 2 & 1 \end{vmatrix} = 2(6-2)-9(1-1)+3(2-6)$$

$$= 2(4) - 9(0) + 3(-4)$$

$$= 8-12$$

$$= -4$$

$$D_z = \begin{vmatrix} + & - & + \\ 2 & -1 & 9 \\ 1 & 1 & 6 \\ 1 & -1 & 2 \end{vmatrix} = 2(2+6)+1(2-6)+9(-1-1)$$

$$= 2(8) + 1(-4) + 9(-2)$$

$$= 16 - 4 - 18$$

$$= -6$$

$$x = \frac{D_x}{D} ; y = \frac{D_y}{D} ; z = \frac{D_z}{D}$$

$$= \frac{-2}{-2} = 1 \quad = \frac{-4}{-2} = 2 \quad = \frac{-6}{-2} = 3$$

SS : {1,2,3}

02. $x + y + z = 6$

$$x - y + z = 2$$

$$2x + y - z = 1$$

$$D = \begin{vmatrix} + & - & + \\ 1 & 1 & 1 \\ 1 & -1 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1(1-1)-1(-1-2)+1(1+2)$$

$$= 1(0) - 1(-3) + 1(3)$$

$$= 3 + 3$$

$$= 6$$

CRAMER'S RULE

$$D_x = \begin{vmatrix} + & - & + \\ 6 & 1 & 1 \\ 2 & -1 & 1 \\ 1 & 1 & -1 \end{vmatrix} = 6(1-1)-1(-2-1)+1(2+1)$$

$$= 6(0) - 1(-3) + 1(3)$$

$$= 3 + 3$$

$$= 6$$

$$D_y = \begin{vmatrix} + & - & + \\ 1 & 6 & 1 \\ 1 & 2 & 1 \\ 2 & 1 & -1 \end{vmatrix} = 1(-2-1)-6(-1-2)+1(1-4)$$

$$= 1(-3) - 6(-3) + 1(-3)$$

$$= -3 + 18 - 3$$

$$= 12$$

$$D_z = \begin{vmatrix} + & - & + \\ 1 & 1 & 6 \\ 1 & -1 & 2 \\ 2 & 1 & 1 \end{vmatrix} = 1(-1-2)-1(1-4)+6(1+2)$$

$$= 1(-3) - 1(-3) + 6(3)$$

$$= -3 + 3 + 18$$

$$= 18$$

$$x = \frac{D_x}{D} ; y = \frac{D_y}{D} ; z = \frac{D_z}{D}$$

$$= \frac{6}{6} = 1 \quad = \frac{12}{6} = 2 \quad = \frac{18}{6} = 3$$

SS : {1,2,3}

03. $x - y + z = 4$

$$2x + y - 3z = 0$$

$$x + y + z = 2$$

$$D = \begin{vmatrix} + & - & + \\ 1 & -1 & 1 \\ 2 & 1 & -3 \\ 1 & 1 & 1 \end{vmatrix} = 1(1+3)+1(2+3)+1(2-1)$$

$$= 1(4) + 1(5) + 1(1)$$

$$= 4 + 5 + 1$$

$$= 10$$

$$D_x = \begin{vmatrix} + & - & + \\ 4 & -1 & 1 \\ 0 & 1 & -3 \\ 2 & 1 & 1 \end{vmatrix} = 4(1+3)+1(0+6)+1(0-2)$$

$$= 4(4) + 1(6) + 1(-2)$$

$$= 16 + 6 - 2$$

$$= 20$$

$$D_y = \begin{vmatrix} + & - & + \\ 1 & 4 & 1 \\ 2 & 0 & -3 \\ 1 & 2 & 1 \end{vmatrix} = 1(0+6)-4(2+3)+1(4-0)$$

$$= 1(6) - 4(5) + 1(4)$$

$$= 6 - 20 + 4$$

$$= -10$$

$$\begin{aligned}
 Dz &= \begin{vmatrix} 1 & -1 & 4 \\ 2 & 1 & 0 \\ 1 & 1 & 2 \end{vmatrix} = 1(2-0) + 1(4-0) + 4(2-1) \\
 &= 1(2) + 1(4) + 4(1) \\
 &= 2 + 4 + 4 \\
 &= 10
 \end{aligned}$$

$$\begin{aligned}
 x &= \frac{Dx}{D} ; y = \frac{Dy}{D} ; z = \frac{Dz}{D} \\
 &= \frac{20}{10} = 2 \quad = \frac{-10}{10} = -1 \quad = \frac{10}{10} = 1
 \end{aligned}$$

$$SS : \{2, -1, 1\}$$

04. $x + y + z = 2$
 $x + 2y + z = 1$
 $5x + y + z = 6$

$$\begin{aligned}
 D &= \begin{vmatrix} 1 & 1 & 1 \\ 1 & 2 & 1 \\ 5 & 1 & 1 \end{vmatrix} = 1(2-1) - 1(1-5) + 1(1-10) \\
 &= 1(1) - 1(-4) + 1(-9) \\
 &= 1 + 4 - 9 \\
 &= -4
 \end{aligned}$$

$$\begin{aligned}
 Dx &= \begin{vmatrix} 2 & 1 & 1 \\ 1 & 2 & 1 \\ 6 & 1 & 1 \end{vmatrix} = 2(2-1) - 1(1-6) + 1(1-12) \\
 &= 2(1) - 1(-5) + 1(-11) \\
 &= 2 + 5 - 11 \\
 &= -4
 \end{aligned}$$

$$\begin{aligned}
 Dy &= \begin{vmatrix} 1 & 2 & 1 \\ 1 & 1 & 1 \\ 5 & 6 & 1 \end{vmatrix} = 1(1-6) - 2(1-5) + 1(6-5) \\
 &= 1(-5) - 2(-4) + 1(1) \\
 &= -5 + 8 + 1 \\
 &= 4
 \end{aligned}$$

$$\begin{aligned}
 Dz &= \begin{vmatrix} 1 & 1 & 2 \\ 1 & 2 & 1 \\ 5 & 1 & 6 \end{vmatrix} = 1(12-1) - 1(6-5) + 2(1-10) \\
 &= 1(11) - 1(1) + 2(-9) \\
 &= 11 - 1 - 18 \\
 &= -8
 \end{aligned}$$

$$\begin{aligned}
 x &= \frac{Dx}{D} ; y = \frac{Dy}{D} ; z = \frac{Dz}{D} \\
 &= \frac{-4}{-4} = 1 \quad = \frac{4}{-4} = -1 \quad = \frac{-8}{-4} = 2
 \end{aligned}$$

$$SS : \{1, -1, 2\}$$

05. $\frac{1}{x} + \frac{1}{y} + \frac{1}{z} = 2$
 $\frac{1}{x} - \frac{2}{y} + \frac{1}{z} = 3$
 $\frac{2}{x} - \frac{1}{y} + \frac{3}{z} = -1$

$$\text{Put } \frac{1}{x} = a ; \frac{1}{y} = b ; \frac{1}{z} = c$$

$$a + b + c = 2$$

$$a - 2b + c = 3$$

$$2a - b + 3c = -1$$

$$\begin{aligned}
 D &= \begin{vmatrix} + & - & + \\ 1 & 1 & 1 \\ 1 & -2 & 1 \\ 2 & -1 & 3 \end{vmatrix} = 1(-6+1) - 1(3-2) + 1(-1+4) \\
 &= 1(-5) - 1(1) + 1(3) \\
 &= -5 - 1 + 3 \\
 &= -3
 \end{aligned}$$

$$\begin{aligned}
 Da &= \begin{vmatrix} + & - & + \\ 2 & 1 & 1 \\ 3 & -2 & 1 \\ -1 & -1 & 3 \end{vmatrix} = 2(-6+1) - 1(9+1) + 1(-3-2) \\
 &= 2(-5) - 1(10) + 1(-5) \\
 &= -10 - 10 - 5 \\
 &= -25
 \end{aligned}$$

$$\begin{aligned}
 Db &= \begin{vmatrix} + & - & + \\ 1 & 2 & 1 \\ 1 & 3 & 1 \\ 2 & -1 & 3 \end{vmatrix} = 1(9+1) - 2(3-2) + 1(-1-6) \\
 &= 1(10) - 2(1) + 1(-7) \\
 &= 10 - 2 - 7 \\
 &= 1
 \end{aligned}$$

$$\begin{aligned}
 Dc &= \begin{vmatrix} + & - & + \\ 1 & 1 & 2 \\ 1 & -2 & 3 \\ 2 & -1 & -1 \end{vmatrix} = 1(2+3) - 1(-1-6) + 2(-1+4) \\
 &= 1(5) - 1(-7) + 2(3) \\
 &= 5 + 7 + 6 \\
 &= 18
 \end{aligned}$$

$$\begin{aligned}
 a &= \frac{Da}{D} ; b = \frac{Db}{D} ; c = \frac{Dc}{D} \\
 &= \frac{-25}{-3} = \frac{25}{3} \quad = \frac{1}{-3} = -\frac{1}{3} \quad = \frac{18}{-3} = -6
 \end{aligned}$$

RESUBS.

$$\frac{1}{x} = \frac{25}{3} ; \frac{1}{y} = -\frac{1}{3} ; \frac{1}{z} = -6$$

$$x = \frac{3}{25} ; y = -3 ; z = -\frac{1}{6}$$

**WITHOUT EXPANSION
SHOW THAT**

08. sum of 3 numbers is 2 . If twice the second number is added to sum of first and third we get 1 . On adding sum of second and third number to 5 times the first number we get 6 . Find the three numbers

$$x + y + z = 2$$

$$x + 2y + z = 1$$

$$5x + y + z = 6 \quad \text{REFER SOLN OF (04)}$$

$$\mathbf{01.} \quad \begin{vmatrix} 1 & 1 & x \\ 1 & x & x^2 \\ 1 & x^2 & x^3 \end{vmatrix} = 0$$

LHS

Taking 'x' common from C₁

$$= x \begin{vmatrix} 1 & 1 & 1 \\ 1 & x & x \\ 1 & x^2 & x^2 \end{vmatrix}$$

= x(0) C₁ & C₂ are identical

$$= 0$$

$$\mathbf{02.} \quad \begin{vmatrix} x+a & x+b & x+c \\ y+a & y+b & y+c \\ z+a & z+b & z+c \end{vmatrix} = 0$$

LHS

C₁ - C₂ , C₂ - C₃

$$= \begin{vmatrix} a-b & b-c & x+c \\ a-b & b-c & y+c \\ a-b & b-c & z+c \end{vmatrix}$$

Taking 'a - b' common from C₁ & 'b - c' common from C₂

$$= (a-b)(b-c) \begin{vmatrix} 1 & 1 & x+c \\ 1 & 1 & y+c \\ 1 & 1 & z+c \end{vmatrix}$$

$$= (a-b)(b-c) (0)$$

..... C₁ & C₂ are identical

$$= 0$$

SOLUTION - QSET 2

$$03. \begin{vmatrix} x-y & x+y & x \\ z-x & z+x & z \\ y-z & y+z & y \end{vmatrix} = 0$$

LHS

C1 + C2

$$= \begin{vmatrix} x-y & x+y & x \\ z-x & z+x & z \\ y-z & y+z & y \end{vmatrix}$$

$$= \begin{vmatrix} 2x & x+y & x \\ 2z & z+x & z \\ 2y & y+z & y \end{vmatrix}$$

Taking '2' common from C1

$$= 2 \begin{vmatrix} x & x+y & x \\ z & z+x & z \\ y & y+z & y \end{vmatrix}$$

= 2(0) C1 & C3 are identical

= 0

$$04. \begin{vmatrix} a-b & b-c & c-a \\ b-c & c-a & a-b \\ c-a & a-b & b-c \end{vmatrix} = 0$$

LHS

C1 + C2

$$= \begin{vmatrix} a-c & b-c & c-a \\ b-a & c-a & a-b \\ c-b & a-b & b-c \end{vmatrix}$$

Taking (-) common from C1

$$= - \begin{vmatrix} c-a & b-c & c-a \\ a-b & c-a & a-b \\ b-c & a-b & b-c \end{vmatrix}$$

= 0 C1 & C3 are identical

$$05. \begin{vmatrix} 1 & a & b+c \\ 1 & b & c+a \\ 1 & c & a+b \end{vmatrix} = 0$$

LHS

C3 + C2

$$= \begin{vmatrix} 1 & a & a+b+c \\ 1 & b & a+b+c \\ 1 & c & a+b+c \end{vmatrix}$$

taking (a + b + c) common from C3

$$= \begin{vmatrix} 1 & a & a+b+c \\ 1 & b & a+b+c \\ 1 & c & a+b+c \end{vmatrix}$$

$$= (a+b+c) \begin{vmatrix} 1 & a & 1 \\ 1 & b & 1 \\ 1 & c & 1 \end{vmatrix}$$

= (a + b + c) (0) C1 & C3 are identical

$$06. \begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ y+z & z+x & x+y \end{vmatrix} = 0$$

LHS

R3 + R2

$$= \begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ x+y+z & x+y+z & x+y+z \end{vmatrix}$$

taking (x + y + z) common from R3

$$= (x+y+z) \begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ 1 & 1 & 1 \end{vmatrix}$$

= (x + y + z) (0) R1 & R3 are identical

$$07. \begin{vmatrix} 1 & xy & z(x+y) \\ 1 & yz & x(y+z) \\ 1 & zx & y(z+x) \end{vmatrix} = 0$$

LHS

$$= \begin{vmatrix} 1 & xy & xz + yz \\ 1 & yz & xy + xz \\ 1 & zx & yz + xy \end{vmatrix}$$

$$= \frac{\cancel{pqr}}{\cancel{pqr}} \begin{vmatrix} lp & mq & nr \\ p^2 & q^2 & r^2 \\ 1 & 1 & 1 \end{vmatrix} = \text{RHS}$$

C₃ + C₂

$$= \begin{vmatrix} 1 & xy & xy + yz + zx \\ 1 & yz & xy + yz + zx \\ 1 & zx & xy + yz + zx \end{vmatrix}$$

taking (xy + yz + zx) common from C₃

$$= (xy + yz + zx) \begin{vmatrix} 1 & xy & 1 \\ 1 & yz & 1 \\ 1 & zx & 1 \end{vmatrix}$$

$$= (xy + yz + zx)(0) \dots \text{C}_1 \text{ \& C}_3 \text{ are identical}$$

$$= 0$$

$$08. \begin{vmatrix} xa & yb & zc \\ a^2 & b^2 & c^2 \\ 1 & 1 & 1 \end{vmatrix} = \begin{vmatrix} x & y & z \\ a & b & c \\ bc & ca & ab \end{vmatrix}$$

RHS

C₁(a) , C₂(b) , C₃(c)

$$= \frac{1}{abc} \begin{vmatrix} xa & yb & zc \\ a^2 & b^2 & c^2 \\ abc & abc & abc \end{vmatrix}$$

taking (abc) common from R₃

$$= \frac{\cancel{abc}}{\cancel{abc}} \begin{vmatrix} xa & yb & zc \\ a^2 & b^2 & c^2 \\ 1 & 1 & 1 \end{vmatrix} = \text{RHS}$$

$$09. \begin{vmatrix} lp & mq & nr \\ p^2 & q^2 & r^2 \\ 1 & 1 & 1 \end{vmatrix} = \begin{vmatrix} l & m & n \\ p & q & r \\ qr & pr & pq \end{vmatrix}$$

RHS

C₁(p) , C₂(q) , C₃(r)

$$= \frac{1}{pqr} \begin{vmatrix} lp & mq & nr \\ p^2 & q^2 & r^2 \\ pqr & pqr & pqr \end{vmatrix}$$

taking (pqr) common from R₃

$$10. \begin{vmatrix} bc & a & a^2 \\ ca & b & b^2 \\ ab & c & c^2 \end{vmatrix} = \begin{vmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{vmatrix}$$

LHS

R₁(a) , R₂(b) , R₃(c)

$$= \frac{1}{abc} \begin{vmatrix} abc & a^2 & a^3 \\ abc & b^2 & b^3 \\ abc & c^2 & c^3 \end{vmatrix}$$

taking (abc) common from C₁

$$= \frac{\cancel{abc}}{\cancel{abc}} \begin{vmatrix} 1 & a^2 & a^3 \\ 1 & b^2 & b^3 \\ 1 & c^2 & c^3 \end{vmatrix} = \text{RHS}$$

$$11. \begin{vmatrix} 1 & x^2 & x^3 \\ 1 & y^2 & y^3 \\ 1 & c^2 & c^3 \end{vmatrix} = \begin{vmatrix} yz & x & x^2 \\ xz & y & y^2 \\ xy & z & z^2 \end{vmatrix}$$

RHS

R₁(x) , R₂(y) , R₃(z)

$$= \frac{1}{xvz} \begin{vmatrix} xyz & x^2 & x^3 \\ xyz & y^2 & y^3 \\ xyz & z^2 & z^3 \end{vmatrix}$$

taking (xyz) common from C₁

$$= \frac{\cancel{xyz}}{\cancel{xyz}} \begin{vmatrix} 1 & x^2 & x^3 \\ 1 & y^2 & y^3 \\ 1 & z^2 & z^3 \end{vmatrix} = \text{RHS}$$

$$12. \begin{vmatrix} x+y & y+z & z+x \\ z+x & x+y & y+z \\ y+z & z+x & x+y \end{vmatrix} = 2 \begin{vmatrix} x & y & z \\ z & x & y \\ y & z & x \end{vmatrix}$$

LHS

C₁ - C₂

$$= \begin{vmatrix} x-z & y+z & z+x \\ z-y & x+y & y+z \\ y-x & z+x & x+y \end{vmatrix}$$

C₁ + C₃

$$= \begin{vmatrix} 2x & y+z & z+x \\ 2z & x+y & y+z \\ 2y & z+x & x+y \end{vmatrix}$$

Taking '2' common from C₁

$$= 2 \begin{vmatrix} x & y+z & z+x \\ z & x+y & y+z \\ y & z+x & x+y \end{vmatrix}$$

C₃ - C₁

$$= 2 \begin{vmatrix} x & y+z & z \\ z & x+y & y \\ y & z+x & x \end{vmatrix}$$

C₂ - C₃

$$= 2 \begin{vmatrix} x & y & z \\ z & x & y \\ y & z & x \end{vmatrix} = \text{RHS}$$

$$13. \begin{vmatrix} a+b & b+c & c+a \\ x+y & y+z & z+x \\ p+q & q+r & r+p \end{vmatrix} = 2 \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix}$$

LHS

C₁ - C₂

$$= \begin{vmatrix} a-c & b+c & c+a \\ x-z & y+z & z+x \\ p-r & q+r & r+p \end{vmatrix}$$

C₁ + C₃

$$= \begin{vmatrix} 2a & b+c & c+a \\ 2x & y+z & z+x \\ 2p & q+r & r+p \end{vmatrix}$$

Taking '2' common from C₁

$$= 2 \begin{vmatrix} a & b+c & c+a \\ x & y+z & z+x \\ p & q+r & r+p \end{vmatrix}$$

C₃ - C₁

$$= 2 \begin{vmatrix} a & b+c & c \\ x & y+z & z \\ p & q+r & r \end{vmatrix}$$

C₂ - C₃

$$= 2 \begin{vmatrix} a & b & c \\ x & y & z \\ p & q & r \end{vmatrix} = \text{RHS}$$

$$14. \begin{vmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{vmatrix} = 0$$

$$\text{let } D = \begin{vmatrix} 0 & a & b \\ -a & 0 & c \\ -b & -c & 0 \end{vmatrix}$$

R ↔ C

$$D = \begin{vmatrix} 0 & -a & -b \\ a & 0 & -c \\ b & c & 0 \end{vmatrix}$$

Taking (-) common from C₁, C₂ & C₃

$$D = (-1)^3 \begin{vmatrix} 0 & -a & -b \\ a & 0 & -c \\ b & c & 0 \end{vmatrix}$$

$$D = -D$$

$$2D = 0 \quad \therefore D = 0$$

$$15. \begin{vmatrix} 0 & x-y & y-z \\ y-x & 0 & z-x \\ z-y & x-z & 0 \end{vmatrix} = 0$$

$$= + \begin{vmatrix} 4 & 1 & 7 \\ 11 & 5 & 2 \\ 10 & 4 & 6 \end{vmatrix}$$

$$\text{let } D = \begin{vmatrix} 0 & x-y & y-z \\ y-x & 0 & z-x \\ z-y & x-z & 0 \end{vmatrix}$$

R₁ + R₂

$$= + \begin{vmatrix} 15 & 6 & 9 \\ 11 & 5 & 2 \\ 10 & 4 & 6 \end{vmatrix}$$

R ↔ C

$$D = \begin{vmatrix} 0 & y-x & z-y \\ x-y & 0 & x-z \\ y-z & z-x & 0 \end{vmatrix}$$

Taking '3' common from R₁

$$= 3 \begin{vmatrix} 5 & 2 & 3 \\ 11 & 5 & 2 \\ 10 & 4 & 6 \end{vmatrix} = \text{RHS}$$

Taking (-) common from C₁, C₂ & C₃

$$D = (-)^3 \begin{vmatrix} 0 & x-y & y-z \\ y-x & 0 & z-x \\ z-y & x-z & 0 \end{vmatrix}$$

$$D = - D$$

$$2D = 0 \quad \therefore D = 0$$

$$16. \begin{vmatrix} 11 & 4 & 10 \\ 2 & 7 & 6 \\ 5 & 1 & 4 \end{vmatrix} = 3 \begin{vmatrix} 5 & 2 & 3 \\ 11 & 5 & 2 \\ 10 & 4 & 6 \end{vmatrix}$$

$$\text{LHS} = \begin{vmatrix} 11 & 4 & 10 \\ 2 & 7 & 6 \\ 5 & 1 & 4 \end{vmatrix}$$

R ↔ C

$$= \begin{vmatrix} 11 & 2 & 5 \\ 4 & 7 & 1 \\ 10 & 6 & 4 \end{vmatrix}$$

C₂ ↔ C₃

$$= - \begin{vmatrix} 11 & 5 & 2 \\ 4 & 1 & 7 \\ 10 & 4 & 6 \end{vmatrix}$$

R₁ ↔ R₂

PROPERTIES OF DETERMINANTS

Q SET - 3

**WITHOUT EXPANDING
AS FAR AS POSSIBLE**

$$01. \begin{vmatrix} a^2+2a & 2a+1 & 1 \\ 2a+1 & a+2 & 1 \\ 3 & 3 & 1 \end{vmatrix} = (a-1)^3$$

R₁ - R₂ , R₂ - R₃

$$= \begin{vmatrix} a^2-1 & a-1 & 0 \\ 2a-2 & a-1 & 0 \\ 3 & 3 & 1 \end{vmatrix}$$

Taking (a - 1) common from R₁ & R₂

$$= (a-1)^2 \begin{vmatrix} a+1 & 1 & 0 \\ 2 & 1 & 0 \\ 3 & 3 & 1 \end{vmatrix}$$

R₁ - R₂

$$= (a-1)^2 \begin{vmatrix} a-1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 3 & 1 \end{vmatrix}$$

Taking (a - 1) common from R₁

$$= (a-1)^3 \begin{vmatrix} 1 & 0 & 0 \\ 2 & 1 & 0 \\ 3 & 3 & 1 \end{vmatrix}$$

Expanding the determinant

$$= (a-1)^3 [1(1-0)]$$

$$= (a-1)^3$$

$$02. \begin{vmatrix} 1 & x & x^2 \\ 1 & y & y^2 \\ 1 & z & z^2 \end{vmatrix} = (x-y)(y-z)(z-x)$$

R₁ - R₂ , R₂ - R₃

$$= \begin{vmatrix} 0 & x-y & x^2-y^2 \\ 0 & y-z & y^2-z^2 \\ 1 & z & z^2 \end{vmatrix}$$

**Taking (x - y) common from R₁ & (y - z)
common from R₂**

$$= (x-y)(y-z) \begin{vmatrix} 0 & 1 & x+y \\ 0 & 1 & y+z \\ 1 & z & z^2 \end{vmatrix}$$

R₁ - R₂

$$= (x-y)(y-z) \begin{vmatrix} 0 & 0 & x-z \\ 0 & 1 & y+z \\ 1 & z & z^2 \end{vmatrix}$$

Taking (x - z) common from R₁

$$= (x-y)(y-z)(x-z) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & y+z \\ 1 & z & z^2 \end{vmatrix}$$

Expanding the determinant

$$= (x-y)(y-z)(x-z) [1(0-1)]$$

$$= (x-y)(y-z)(x-z) (-1)$$

$$= (x-y)(y-z)(z-x)$$

$$03. \begin{vmatrix} 1 & a & bc \\ 1 & b & ca \\ 1 & c & ab \end{vmatrix} = (a-b)(b-c)(c-a)$$

$$= abc \begin{vmatrix} 1 & 1 & 1 \\ a & b & c \\ a^2 & b^2 & c^2 \end{vmatrix}$$

R₁ - R₂ , R₂ - R₃

$$= \begin{vmatrix} 0 & a-b & bc-ca \\ 0 & b-c & ca-ab \\ 1 & c & ab \end{vmatrix}$$

$$= \begin{vmatrix} 0 & a-b & c(b-a) \\ 0 & b-c & a(c-b) \\ 1 & c & ab \end{vmatrix}$$

Taking (a - b) common from R₁ & (b - c) common from R₂

$$= (a-b)(b-c) \begin{vmatrix} 0 & 1 & -c \\ 0 & 1 & -a \\ 1 & c & ab \end{vmatrix}$$

R₁ - R₂

$$= (a-b)(b-c) \begin{vmatrix} 0 & 0 & a-c \\ 0 & 1 & -a \\ 1 & c & ab \end{vmatrix}$$

Taking (a - c) common from R₁

$$= (a-b)(b-c)(a-c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & -a \\ 1 & c & ab \end{vmatrix}$$

$$= (a-b)(b-c)(a-c) [1(0-1)]$$

$$= (a-b)(b-c)(a-c)(-1)$$

$$= (a-b)(b-c)(c-a)$$

$$04. \begin{vmatrix} a & b & c \\ a^2 & b^2 & c^2 \\ a^3 & b^3 & c^3 \end{vmatrix} = abc(a-b)(b-c)(c-a)$$

taking 'a' , 'b' & 'c' common from C₁ , C₂ & C₃ respectively

C₁ - C₂ , C₂ - C₃

$$= abc \begin{vmatrix} 0 & 0 & 1 \\ a-b & b-c & c \\ a^2-b^2 & b^2-c^2 & c^2 \end{vmatrix}$$

Taking (a - b) common from C₁ & (b - c) common from C₂

$$= abc(a-b)(b-c) \begin{vmatrix} 0 & 0 & 1 \\ 1 & 1 & c \\ a+b & b+c & c^2 \end{vmatrix}$$

C₁ - C₂

$$= abc(a-b)(b-c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & c \\ a-c & b+c & c^2 \end{vmatrix}$$

Taking (a - c) common from C₁

$$= abc(a-b)(b-c)(a-c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & c \\ 1 & b+c & c^2 \end{vmatrix}$$

Expanding the determinant

$$= abc(a-b)(b-c)(a-c) [1(0-1)]$$

$$= abc(a-b)(b-c)(a-c) (-1)$$

$$= abc(a-b)(b-c)(c-a)$$

$$05. \begin{vmatrix} x & p & q \\ p & x & q \\ p & q & x \end{vmatrix} = (x-p)(x-q)(x+p+q)$$

R₁ - R₂ , R₂ - R₃

$$= \begin{vmatrix} x-p & p-x & 0 \\ 0 & x-q & q-x \\ p & q & x \end{vmatrix}$$

Taking (x - p) common from R₁ & (x - q) common from R₂

$$= (x-p)(x-q) \begin{vmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ p & q & x \end{vmatrix}$$

Expanding the determinant

$$= (x-p)(x-q) [1(x+q) + 1(0+p)]$$

$$= (x-p)(x-q)(x+p+q)$$

$$06. \begin{vmatrix} a & b+c & a^2 \\ b & c+a & b^2 \\ c & a+b & c^2 \end{vmatrix} = -(a-b)(b-c)(c-a)(a+b+c)$$

C₁ - C₂

$$= \begin{vmatrix} a+b+c & b+c & a^2 \\ a+b+c & c+a & b^2 \\ a+b+c & a+b & c^2 \end{vmatrix}$$

Taking (a - c) common from C₁

$$= (a+b+c) \begin{vmatrix} 1 & b+c & a^2 \\ 1 & c+a & b^2 \\ 1 & a+b & c^2 \end{vmatrix}$$

R₁ - R₂ , R₂ - R₃

$$= (a+b+c) \begin{vmatrix} 0 & b-a & a^2-b^2 \\ 0 & c-b & b^2-c^2 \\ 1 & a+b & c^2 \end{vmatrix}$$

Taking (a - b) common from R₁ & (b - c) common from R₂

$$= (a+b+c)a-b)(b-c) \begin{vmatrix} 0 & -1 & a+b \\ 0 & -1 & b+c \\ 1 & a+b & c^2 \end{vmatrix}$$

R₁ - R₂

$$= (a+b+c)a-b)(b-c) \begin{vmatrix} 0 & 0 & a-c \\ 0 & -1 & b+c \\ 1 & a+b & c^2 \end{vmatrix}$$

Taking (a - c) common from R₁

$$= (a+b+c)a-b)(b-c)(a-c) \begin{vmatrix} 0 & 0 & 1 \\ 0 & -1 & b+c \\ 1 & a+b & c^2 \end{vmatrix}$$

Expanding the determinant

$$= (a+b+c)(a-b)(b-c)(a-c) [1(0+1)]$$

$$= (a+b+c)(a-b)(b-c)(a-c) (1)$$

$$= -(a+b+c)(a-b)(b-c)(c-a)$$

$$07. \begin{vmatrix} a+b+2c & a & b \\ c & b+c+2a & b \\ c & a & c+a+2b \end{vmatrix} = 2(a+b+c)^3$$

R₁ - R₂ , R₂ - R₃

$$= \begin{vmatrix} a+b+c & -a-b-c & 0 \\ 0 & b+c+a & -b-c-a \\ c & a & c+a+2b \end{vmatrix}$$

Taking (a+b+c) common from R₁ & R₂ respectively .

$$= (a+b+c)^2 \begin{vmatrix} 1 & -1 & 0 \\ 0 & 1 & -1 \\ c & a & c+a+2b \end{vmatrix}$$

Expanding the determinant

$$= (a+b+c)^2 [1(c+a+2b+a) + 1(0+c)]$$

$$= (a+b+c)^2 (2a+2b+2c)$$

$$= (a+b+c)^2 2(a+b+c)$$

$$= 2(a+b+c)^3$$

$$08. \begin{vmatrix} 1 & 1 & 1 \\ x & y & z \\ x^3 & y^3 & z^3 \end{vmatrix} = (x-y)(y-z)(z-x)(x+y+z)$$

C₁ - C₂ , C₂ - C₃

$$= \begin{vmatrix} 0 & 0 & 1 \\ x-y & y-z & z \\ x^3-y^3 & y^3-z^3 & z^3 \end{vmatrix}$$

Taking (x - y) common from C₁ & (y - z) common from C₂

$$= (x-y)(y-z) \begin{vmatrix} 0 & 0 & 1 \\ 1 & 1 & z \\ x^2+xy+y^2 & y^2+yz+z^2 & z^3 \end{vmatrix}$$

C₁ - C₂

$$= (x-y)(y-z) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & z \\ x^2-z^2+xy-yz & y^2+yz+z^2 & z^3 \end{vmatrix}$$

$$= (x-y)(y-z) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & z \\ (x-z)(x+z)+y(x-z) & y^2+yz+z^2 & z^3 \end{vmatrix}$$

$$= (x-y)(y-z) \begin{vmatrix} 0 & 0 & 1 \\ 0 & 1 & z \\ (x-z)(x+y+z) & y^2+yz+z^2 & z^3 \end{vmatrix}$$

Expanding the determinant

$$= (x-y)(y-z) \cdot 1(0 - (x-z)(x+y+z))$$

$$= (x-y)(y-z)(z-x)(x+y+z)$$

$$09. \begin{vmatrix} y+z & z & y \\ z & z+x & x \\ y & x & x+y \end{vmatrix} = 4xyz$$

C₁ - C₂

$$= \begin{vmatrix} y & z & y \\ -x & z+x & x \\ y-x & x & x+y \end{vmatrix}$$

C₁ + C₃

$$= \begin{vmatrix} 2y & z & y \\ 0 & z+x & x \\ 2y & x & x+y \end{vmatrix}$$

Taking '2' common from C₁

$$= 2 \begin{vmatrix} y & z & y \\ 0 & z+x & x \\ y & x & x+y \end{vmatrix}$$

C₃ - C₁

$$= 2 \begin{vmatrix} y & z & 0 \\ 0 & z+x & x \\ y & x & x \end{vmatrix}$$

C₂ - C₃

$$= 2 \begin{vmatrix} y & z & 0 \\ 0 & z & x \\ y & 0 & x \end{vmatrix}$$

Taking 'x' , 'y' & z common from C₁,C₂ & C₃ respectively

$$= 2xyz \begin{vmatrix} 1 & 1 & 0 \\ 0 & 1 & 1 \\ 1 & 0 & 1 \end{vmatrix}$$

$$= 2xyz [1(1-0) - 1(0-1)]$$

$$= 2xyz(1+1) = 4xyz$$

10. WITHOUT EXPANSION , PROVE

$$\begin{vmatrix} 6 & 1 & 2 \\ 7 & 2 & 3 \\ 2 & 3 & -4 \end{vmatrix} + \begin{vmatrix} 2 & 4 & 7 \\ 3 & 4 & 2 \\ -4 & 3 & 3 \end{vmatrix} = 5 \begin{vmatrix} 2 & 2 & 7 \\ 3 & 1 & 2 \\ -4 & 1 & 3 \end{vmatrix}$$

LHS

$$= \begin{vmatrix} 6 & 1 & 2 \\ 7 & 2 & 3 \\ 2 & 3 & -4 \end{vmatrix} + \begin{vmatrix} 2 & 4 & 7 \\ 3 & 4 & 2 \\ -4 & 3 & 3 \end{vmatrix}$$

R ↔ C (ONLY ON THE FIRST DETERMINANT)

$$= \begin{vmatrix} 6 & 7 & 2 \\ 1 & 2 & 3 \\ 2 & 3 & -4 \end{vmatrix} + \begin{vmatrix} 2 & 4 & 7 \\ 3 & 4 & 2 \\ -4 & 3 & 3 \end{vmatrix}$$

C₂ ↔ C₃ (ONLY ON THE FIRST DETERMINANT)

$$= - \begin{vmatrix} 6 & 2 & 7 \\ 1 & 3 & 2 \\ 2 & -4 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 4 & 7 \\ 3 & 4 & 2 \\ -4 & 3 & 3 \end{vmatrix}$$

C₁ ↔ C₂ (ONLY ON THE FIRST DETERMINANT)

$$= \begin{vmatrix} 2 & 6 & 7 \\ 3 & 1 & 2 \\ -4 & 2 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 4 & 7 \\ 3 & 4 & 2 \\ -4 & 3 & 3 \end{vmatrix}$$

$$= \begin{vmatrix} 2 & 6+4 & 7 \\ 3 & 1+4 & 2 \\ -4 & 2+3 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 4 & 7 \\ 3 & 4 & 2 \\ -4 & 3 & 3 \end{vmatrix}$$

$$= \begin{vmatrix} 2 & 10 & 7 \\ 3 & 5 & 2 \\ -4 & 5 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 4 & 7 \\ 3 & 4 & 2 \\ -4 & 3 & 3 \end{vmatrix}$$

taking '5' common from C₂

$$= 5 \begin{vmatrix} 2 & 2 & 7 \\ 3 & 1 & 2 \\ -4 & 1 & 3 \end{vmatrix} = \text{RHS}$$

11. WITHOUT EXPANSION , PROVE

$$\begin{vmatrix} 2 & -3 & 4 \\ 5 & -6 & 2 \\ -3 & 1 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 5 & -3 \\ 5 & 8 & 5 \\ 4 & 2 & 3 \end{vmatrix} = -2 \begin{vmatrix} 2 & 4 & 1 \\ 5 & 2 & 1 \\ -3 & 3 & 3 \end{vmatrix}$$

LHS

$$= \begin{vmatrix} 2 & -3 & 4 \\ 5 & -6 & 2 \\ -3 & 1 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 5 & -3 \\ 5 & 8 & 5 \\ 4 & 2 & 3 \end{vmatrix}$$

R ↔ C (ONLY ON THE SECOND DETERMINANT)

$$= \begin{vmatrix} 2 & -3 & 4 \\ 5 & -6 & 2 \\ -3 & 1 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 5 & 4 \\ 5 & 8 & 2 \\ -3 & 5 & 3 \end{vmatrix}$$

$$= \begin{vmatrix} 2 & -3+5 & 4 \\ 5 & -6+8 & 2 \\ -3 & 1+5 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 5 & 4 \\ 5 & 8 & 2 \\ -3 & 5 & 3 \end{vmatrix}$$

$$= \begin{vmatrix} 2 & 2 & 4 \\ 5 & 2 & 2 \\ -3 & 6 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 5 & 4 \\ 5 & 8 & 2 \\ -3 & 5 & 3 \end{vmatrix}$$

taking '2' common from C₂

$$= 2 \begin{vmatrix} 2 & 1 & 4 \\ 5 & 1 & 2 \\ -3 & 3 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 5 & 4 \\ 5 & 8 & 2 \\ -3 & 5 & 3 \end{vmatrix}$$

C₂ ↔ C₃ (ONLY ON THE SECOND DETERMINANT)

$$= -2 \begin{vmatrix} 2 & 4 & 1 \\ 5 & 2 & 1 \\ -3 & 3 & 3 \end{vmatrix} + \begin{vmatrix} 2 & 5 & 4 \\ 5 & 8 & 2 \\ -3 & 5 & 3 \end{vmatrix}$$

= RHS

SOLUTION - QSET 4**CONSISTENCY OF EQUATIONS**

01. $x + y = 3$; $5x + 6y = 17$; $2x - 3y = k$
are consistent . Find k

$$x + y = 3$$

$$5x + 6y = 17$$

$$2x - 3y = k \quad \text{are consistent}$$

Hence

$$\begin{vmatrix} + & - & + \\ 1 & 1 & 3 \\ 5 & 6 & 17 \\ 2 & -3 & k \end{vmatrix} = 0$$

$$1(6k + 51) - 1(5k - 34) + 3(-15 - 12) = 0$$

$$6k + 51 - 5k + 34 + 3(-27) = 0$$

$$k + 85 - 81 = 0$$

$$k + 4 = 0$$

$$k = -4$$

02. $3x + y = 2$; $kx + 2y = 3$; $2x - y = -3$
are consistent . Find k

$$3x + y = 2$$

$$kx + 2y = 3$$

$$2x - y = -3 \quad \text{are consistent}$$

Hence

$$\begin{vmatrix} + & - & + \\ 3 & 1 & 2 \\ k & 2 & 3 \\ 2 & -1 & -3 \end{vmatrix} = 0$$

$$3(-6 + 3) - 1(-3k - 6) + 2(-k - 4) = 0$$

$$3(-3) + 3k + 6 - 2k - 8 = 0$$

$$-9 + k - 2 = 0$$

$$-11 + k = 0$$

$$k = 11$$

03. $x + 3y + 2 = 0$; $4y + 2x = k$; $x - 2y = 3k$.
are consistent . Find k

$$x + 3y = -2$$

$$2x + 4y = k$$

$$x - 2y = 3k \quad \text{are consistent}$$

Hence

$$\begin{vmatrix} + & - & + \\ 1 & 3 & -2 \\ 2 & 4 & k \\ 1 & -2 & 3k \end{vmatrix} = 0$$

$$1(12k + 2k) - 3(6k - k) - 2(-4 - 4) = 0$$

$$1(14k) - 18k + 3k - 2(-8) = 0$$

$$14k - 18k + 3k + 16 = 0$$

$$-k + 16 = 0$$

$$k = 16$$

04. $2x - y + 3 = 0$; $7x - 2y + 2 = 0$;
 $kx - y - 1 = 0$ are consistent . Find k & SS

$$2x - y = -3$$

$$7x - 2y = -2$$

$$kx - y = 1 \quad \text{are consistent}$$

Hence

$$\begin{vmatrix} + & - & + \\ 2 & -1 & -3 \\ 7 & -2 & -2 \\ k & -1 & 1 \end{vmatrix} = 0$$

$$2(-2 - 2) + 1(7 + 2k) - 3(-7 + 2k) = 0$$

$$2(-4) + 7 + 2k + 21 - 6k = 0$$

$$-8 + 28 - 4k = 0$$

$$20 - 4k = 0$$

$$20 = 4k$$

$$k = 5$$

FOR SOLUTION SET

$$D = \begin{vmatrix} 2 & -1 \\ 7 & -2 \end{vmatrix} = -4 + 7 = 3$$

$$D_x = \begin{vmatrix} -3 & -1 \\ -2 & -2 \end{vmatrix} = 6 - 2 = 4$$

$$D_y = \begin{vmatrix} 2 & -3 \\ 7 & -2 \end{vmatrix} = -4 + 21 = 17$$

$$x = \frac{D_x}{D} = \frac{4}{3} ; y = \frac{D_y}{D} = \frac{17}{3} \quad \text{SS} : \left\{ \frac{4}{3}, \frac{17}{3} \right\}$$

05. $3x + y = 2$; $kx + 2y = 3$; $2x - y = -3$.

Find k if the lines are concurrent . Hence find the point of concurrency

$$3x + y = 2$$

$$kx + 2y = 3$$

$$2x - y = -3 \quad \text{are concurrent}$$

Hence

$$\begin{vmatrix} + & - & + \\ 3 & 1 & 2 \\ k & 2 & 3 \\ 2 & -1 & -3 \end{vmatrix} = 0$$

$$3(-6 + 3) - 1(-3k - 6) + 2(-k - 4) = 0$$

$$3(-3) + 3k + 6 - 2k - 8 = 0$$

$$-9 + k - 2 = 0$$

$$-11 + k = 0$$

$$k = 11$$

FOR SOLUTION SET

$$3x + y = 2$$

$$2x - y = -3$$

$$D = \begin{vmatrix} 3 & 1 \\ 2 & -1 \end{vmatrix} = -3 - 2 = -5$$

$$Dx = \begin{vmatrix} 2 & 1 \\ -3 & -1 \end{vmatrix} = -2 + 3 = 1$$

$$Dy = \begin{vmatrix} 3 & 2 \\ 2 & -3 \end{vmatrix} = -9 - 4 = -13$$

$$x = \frac{Dx}{D} = \frac{-1}{-5}; \quad y = \frac{Dy}{D} = \frac{-13}{-5} \quad \text{SS} : \left\{ \frac{-1}{5}, \frac{13}{5} \right\}$$

06. $kx + 3y + 4 = 0$; $x + ky + 3 = 0$;

$$3x + 4y + 5 = 0 \quad \text{are consistent .}$$

Find k . Hence find common solution set for smallest value of k

$$kx + 3y = -4$$

$$x + ky = -3$$

$$3x + 4y = -5 \quad \text{are consistent}$$

$$\begin{vmatrix} k & 3 & -4 \\ 1 & k & -3 \\ 3 & 4 & -5 \end{vmatrix} = 0$$

taking - common from C_1

$$\begin{vmatrix} + & - & + \\ k & 3 & 4 \\ 1 & k & 3 \\ 3 & 4 & 5 \end{vmatrix} = 0$$

$$k(5k - 12) - 3(5 - 9) + 4(4 - 3k) = 0$$

$$5k^2 - 12k - 3(-4) + 16 - 12k = 0$$

$$5k^2 - 12k + 12 + 16 - 12k = 0$$

$$5k^2 - 24k + 28 = 0$$

$$5k^2 - 10k - 14k + 28 = 0$$

$$5k(k - 2) - 14(k - 2) = 0$$

$$k = 14/5, \quad k = 2$$

SOLUTION SET FOR $k = 2$

$$2x + 3y = -4$$

$$x + 2y = -3$$

FOR SOLUTION SET

$$D = \begin{vmatrix} 2 & 3 \\ 1 & 2 \end{vmatrix} = 4 - 3 = 1$$

$$Dx = \begin{vmatrix} -4 & 3 \\ -3 & 2 \end{vmatrix} = -8 + 9 = 1$$

$$Dy = \begin{vmatrix} 2 & -4 \\ 1 & -3 \end{vmatrix} = -6 + 4 = -2$$

$$x = \frac{Dx}{D} = \frac{1}{1}; \quad y = \frac{Dy}{D} = \frac{-2}{1} \quad \text{SS} : \{1, -2\}$$

07. if the equations $ax + by + c = 0$;
 $cx + ay + b = 0$; and $bx + cy + a = 0$
 are consistent in x and y then show that
 $a^3 + b^3 + c^3 = 3abc$

$$\begin{aligned} ax + by &= -c \\ cx + ay &= -b \\ bx + cy &= -a \end{aligned} \text{ are consistent}$$

Hence

$$\begin{vmatrix} a & b & -c \\ c & a & -b \\ b & c & -a \end{vmatrix} = 0$$

taking '- ' common from C₁

$$\begin{vmatrix} + & - & + \\ a & b & c \\ c & a & b \\ b & c & a \end{vmatrix} = 0$$

Expanding the determinant

$$a(a^2 - bc) - b(ac - b^2) + c(c^2 - ab) = 0$$

$$a^3 - abc - abc + b^3 + c^3 - abc = 0$$

$$a^3 + b^3 + c^3 - 3abc = 0$$

$$a^3 + b^3 + c^3 = 3abc \dots\dots \text{PROVED}$$

08. if $l = \frac{y-1}{x}$; $m = \frac{1-x}{y}$; $n = x-y$

are consistent in x and y then prove :

$$l + m + n + lmn = 0$$

$$lx - y = -1$$

$$x + my = 1$$

$x - y = n$ are consistent , hence

$$\begin{vmatrix} + & - & + \\ l & -1 & -1 \\ 1 & m & 1 \\ 1 & -1 & n \end{vmatrix} = 0$$

Expanding the determinant

$$l(mn + 1) + 1(n - 1) - 1(-1 - m) = 0$$

$$lmn + l + n - 1 + 1 + m = 0$$

$$lmn + l + m + n = 0$$

SOLUTION - QSET 5

Q1. Find area of triangle with vertices as

01. (4,5) ; (0,7) ; (-1,1)

$$\begin{aligned} A &= \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \\ &= \frac{1}{2} \begin{vmatrix} 4 & 5 & 1 \\ 0 & 7 & 1 \\ -1 & 1 & 1 \end{vmatrix} \\ &= \frac{1}{2} [4(7-1) - 5(0+1) + 1(0+7)] \\ &= \frac{1}{2} [24 - 5 + 7] \\ &= \frac{1}{2} (26) = 13 \text{ sq. units} \end{aligned}$$

02. (3,6) ; (-1,3) ; (2,-1)

$$\begin{aligned} A &= \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \\ &= \frac{1}{2} \begin{vmatrix} 3 & 6 & 1 \\ -1 & 3 & 1 \\ 2 & -1 & 1 \end{vmatrix} \\ &= \frac{1}{2} [3(3+1) - 6(-1-2) + 1(1-6)] \\ &= \frac{1}{2} [12 + 18 - 5] \\ &= \frac{1}{2} (25) = \frac{25}{2} \text{ sq. units} \end{aligned}$$

03. (1,1) , (3,7) , (10,8)

$$\begin{aligned} A &= \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \\ &= \frac{1}{2} \begin{vmatrix} 1 & 1 & 1 \\ 3 & 7 & 1 \\ 10 & 8 & 1 \end{vmatrix} \\ &= \frac{1}{2} [1(7-8) - 1(3-10) + 1(24-70)] \end{aligned}$$

AREA OF TRIANGLE

$$\begin{aligned} &= \frac{1}{2} [-1 + 7 - 46] \\ &= \frac{1}{2} (-40) \\ &= -20 = 20 \text{ sq. units} \end{aligned}$$

Q2.

01. Find k if the area of the triangle whose vertices are A (4 , k) ; B (-5 , -7) ; C(-4 , 1) is 38 sq. units

$$A (\Delta ABC) = 38$$

$$\begin{aligned} \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} &= \pm 38 \\ \begin{vmatrix} 4 & k & 1 \\ -5 & -7 & 1 \\ -4 & 1 & 1 \end{vmatrix} &= \pm 76 \end{aligned}$$

$$4(-7-1) - k(-5+4) + 1(-5-28) = \pm 76$$

$$4(-8) - k(-1) + 1(-33) = \pm 76$$

$$-32 + k - 33 = \pm 76$$

$$-65 + k = \pm 76$$

$$-65 + k = 76 \quad \text{OR} \quad -65 + k = -76$$

$$k = 141$$

$$k = -11$$

02. Find k if the area of the triangle whose vertices are P (k , -4) ; Q (1 , -2) ; R (4 , -5) is $15/2$ sq. units

$$A (\Delta ABC) = 15/2$$

$$\begin{aligned} \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} &= \pm \frac{15}{2} \\ \begin{vmatrix} k & -4 & 1 \\ 1 & -2 & 1 \\ 4 & -5 & 1 \end{vmatrix} &= \pm 15 \end{aligned}$$

$$k(-2+5) + 4(1-4) + 1(-5+8) = \pm 15$$

$$3k - 12 + 3 = \pm 15$$

$$3k - 9 = \pm 15$$

$$3k - 9 = 15 \quad \text{OR} \quad 3k - 9 = -15$$

$$3k = 24 \qquad 3k = -6$$

$$k = 8 \qquad \text{OR} \quad k = -2$$

03. Find k if the area of the triangle whose vertices are P (3, -5) ; Q (-2, k) ; R (1, 4)

is $33/2$ sq. units

$$A(\Delta ABC) = 33/2$$

$$\frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = \pm \frac{33}{2}$$

$$\begin{vmatrix} 3 & -5 & 1 \\ -2 & k & 1 \\ 1 & 4 & 1 \end{vmatrix} = \pm 33$$

$$3(k - 4) + 5(-2 - 1) + 1(-8 - k) = \pm 33$$

$$3k - 12 - 15 - 8 - k = \pm 33$$

$$2k - 35 = \pm 33$$

$$2k - 35 = 33 \quad \text{OR} \quad 2k - 35 = -33$$

$$2k = 68 \qquad 2k = 2$$

$$k = 34 \qquad \text{OR} \quad k = 1$$

Q3. Find area of Quadrilateral whose vertices are A (2, 1) ; B (2, 3) ; C(-2, 2) ; D(-1, 0)

$$A(\Delta ABC) = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

$$= \frac{1}{2} \begin{vmatrix} 2 & 1 & 1 \\ 2 & 3 & 1 \\ -2 & 2 & 1 \end{vmatrix}$$

$$= \frac{1}{2} [2(3-2) - 1(2+2) + 1(4+6)]$$

$$= \frac{1}{2} [2 - 4 + 10]$$

$$= \frac{1}{2} (8) = 4 \text{ sq. units}$$

$$A(\Delta ADC) = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

$$= \frac{1}{2} \begin{vmatrix} 2 & 1 & 1 \\ -1 & 0 & 1 \\ -2 & 2 & 1 \end{vmatrix}$$

$$= \frac{1}{2} [2(0-2) - 1(-1+2) + 1(-2+0)]$$

$$= \frac{1}{2} [-4 - 1 - 2]$$

$$= -\frac{7}{2} = \frac{7}{2} \text{ sq. units}$$

$$A(\square ABCD) = A(\Delta ABC) + A(\Delta ADC)$$

$$= 4 + \frac{7}{2}$$

$$= 15/2 \text{ sq. units}$$

Q4.

Using determinants show that the following set of points are collinear

01. A(3,1) ; B(4,2) ; C(5, 3)

$$A = \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix}$$

$$= \frac{1}{2} \begin{vmatrix} 3 & 1 & 1 \\ 4 & 2 & 1 \\ 5 & 3 & 1 \end{vmatrix}$$

$$= \frac{1}{2} [3(2-3) - 1(4-5) + 1(12-10)]$$

$$= \frac{1}{2} [-3 + 1 + 2]$$

$$= 0$$

Hence the points are collinear

02. A(1,-2) ; B(3,1) ; C(5,4)

$$\begin{aligned}
 A &= \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \\
 &= \frac{1}{2} \begin{vmatrix} 1 & -2 & 1 \\ 3 & 1 & 1 \\ 5 & 4 & 1 \end{vmatrix} \\
 &= \frac{1}{2} [1(1-4) + 2(3-5) + 1(12-5)] \\
 &= \frac{1}{2} [-3 - 4 + 7] \\
 &= 0
 \end{aligned}$$

Hence the points are collinear

03. A(3,7) ; B(4,-3) ; C(5,-13)

$$\begin{aligned}
 A &= \frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} \\
 &= \frac{1}{2} \begin{vmatrix} 3 & 7 & 1 \\ 4 & -3 & 1 \\ 5 & -13 & 1 \end{vmatrix} \\
 &= \frac{1}{2} [3(-3+13) - 7(4-5) + 1(-52+15)] \\
 &= \frac{1}{2} [3(10) - 7(-1) - 1(37)] \\
 &= \frac{1}{2} [30 + 7 - 37] \\
 &= 0
 \end{aligned}$$

Hence the points are collinear

04.

points A(a,0) ; B(0,b) , C(1,1) are collinear .

Prove : $\frac{1}{a} + \frac{1}{b} = 1$

Since points are collinear ;

$$A(\Delta ABC) = 0$$

$$\frac{1}{2} \begin{vmatrix} x_1 & y_1 & 1 \\ x_2 & y_2 & 1 \\ x_3 & y_3 & 1 \end{vmatrix} = 0$$

$$\frac{1}{2} \begin{vmatrix} a & 0 & 1 \\ 0 & b & 1 \\ 1 & 1 & 1 \end{vmatrix} = 0$$

$$a(b-1) - 0(0-1) + 1(0-b) = 0$$

$$ab - a - b = 0$$

$$ab = a + b$$

Dividing throughout by ab

$$1 = \frac{a}{ab} + \frac{b}{ab}$$

$$\frac{1}{a} + \frac{1}{b} = 1 \quad \dots\dots \text{PROVED}$$

05 if points A(a ,b) ;B(c , d) ; C(a - c , b - d) are collinear then prove that ad - bc = 0

Since points are collinear ;

$$A(\Delta ABC) = 0$$

$$\frac{1}{2} \begin{vmatrix} a & b & 1 \\ c & d & 1 \\ a-c & b-d & 1 \end{vmatrix} = 0$$

$$\begin{vmatrix} a & b & 1 \\ c & d & 1 \\ a-c & b-d & 1 \end{vmatrix} = 0$$

$$a(d-b+d) - b(c-a+c) + 1(bc-cd-ad+cd) = 0$$

$$a(2d-b) - b(2c-a) + 1(bc-ad) = 0$$

$$2ad - ab - 2bc + ab + bc - ad = 0$$

$$ad - bc = 0$$