

FYJC - MATHEMATICS & STATISTICS

HIGHLIGHTS

- ✓ *Solution to all questions*
- ✓ *solutions are put in way the student is expected to reproduce in the exam*
- ✓ *taught in the class room the same way as the solution are put up here . That makes the student to easily go through the solution & prepare him/herself when he/she sits back to revise and recall the topic at any given point of time .*
- ✓ *lastly, if student due to some unavoidable reasons , has missed the lecture , will not have to run here and there to update his/her notes .*
- ✓ *however class room lectures are must for easy passage of understanding & learning the minuest details of the given topic*

PAPER - I

CIRCULAR INVERSE TRIGONOMETRIC FN'S

Q SET - 1

$$\sin^{-1}(\sin\theta) = \theta$$

$$\cos^{-1}(\cos\theta) = \theta$$

$$\tan^{-1}(\tan\theta) = \theta$$

$$\sin(\sin^{-1}x) = x$$

$$\cos(\cos^{-1}x) = x$$

$$\tan(\tan^{-1}x) = x$$

$$\operatorname{cosec}^{-1}(1/x) = \sin^{-1}x$$

$$\sec^{-1}(1/x) = \cos^{-1}x$$

$$\cot^{-1}(1/x) = \tan^{-1}x$$

$$\sin^{-1}x + \cos^{-1}x = \pi/2$$

$$\tan^{-1}x + \cot^{-1}x = \pi/2$$

$$\sec^{-1}x + \operatorname{cosec}^{-1}x = \pi/2$$

$$\tan^{-1}x - \tan^{-1}y = \tan^{-1}\left(\frac{x-y}{1+xy}\right)$$

$$\tan^{-1}x + \tan^{-1}y = \tan^{-1}\left(\frac{x+y}{1-xy}\right)$$

HOWEVER if $xy > 1$ THEN

$$\tan^{-1}x + \tan^{-1}y = \pi + \tan^{-1}\left(\frac{x+y}{1-xy}\right)$$

$$01. \tan^{-1}\left(\frac{2}{3}\right) + \tan^{-1}\left(\frac{1}{5}\right) = \frac{\pi}{4}$$

$$02. \tan^{-1}\left(\frac{7}{9}\right) + \tan^{-1}\left(\frac{1}{8}\right) = \frac{\pi}{4}$$

$$03. \tan^{-1}\left(\frac{3}{5}\right) + \tan^{-1}\left(\frac{1}{4}\right) = \frac{\pi}{4}$$

$$04. \cot^{-1}[8] + \cot^{-1}\left[\frac{9}{7}\right] = \frac{\pi}{4}$$

05.

$$\tan^{-1}\left(\frac{1}{2}\right) + \tan^{-1}\left(\frac{1}{5}\right) + \tan^{-1}\left(\frac{1}{8}\right) = \frac{\pi}{4}$$

06.

$$\tan^{-1}\left(\frac{3}{4}\right) + \tan^{-1}\left(\frac{3}{5}\right) - \tan^{-1}\left(\frac{8}{19}\right) = \frac{\pi}{4}$$

07.

$$2\tan^{-1}\left(\frac{1}{3}\right) + \tan^{-1}\left(\frac{1}{4}\right) = \tan^{-1}\left(\frac{16}{13}\right)$$

08.

$$2\tan^{-1}\left(\frac{1}{8}\right) + \tan^{-1}\left(\frac{1}{7}\right) + 2\tan^{-1}\left(\frac{1}{5}\right) = \frac{\pi}{4}$$

Q SET - 2

$$01. \cos^{-1}\left(\frac{4}{5}\right) + \cos^{-1}\left(\frac{12}{13}\right) = \cos^{-1}\left(\frac{33}{65}\right)$$

$$02. \sin^{-1}\left(\frac{3}{5}\right) + \cos^{-1}\left(\frac{12}{13}\right) = \sin^{-1}\left(\frac{56}{65}\right)$$

$$03. \sin^{-1}\left(\frac{3}{5}\right) + \sin^{-1}\left(\frac{8}{17}\right) = \sin^{-1}\left(\frac{77}{85}\right)$$

$$04. \cos^{-1} \left(\frac{12}{13} \right) + \sin^{-1} \left(\frac{5}{13} \right) = \cot^{-1} \left(\frac{119}{120} \right)$$

$$05. 2\sin^{-1} \left(\frac{3}{5} \right) - \tan^{-1} \left(\frac{17}{31} \right) = \frac{\pi}{4}$$

$$06. \sin^{-1} \left(\frac{3}{\sqrt{34}} \right) + \cos^{-1} \left(\frac{4}{\sqrt{17}} \right) = \frac{\pi}{4}$$

$$07. \sin^{-1} \left(\frac{2}{\sqrt{13}} \right) + \cos^{-1} \left(\frac{5}{\sqrt{26}} \right) = \frac{\pi}{4}$$

Q SET - 3

$$01. \tan^{-1} \left(\frac{\sin 2x}{1 + \cos 2x} \right) = x$$

$$02. \cot^{-1} \left(\frac{\sin 2x}{1 - \cos 2x} \right) = x$$

$$03. \tan^{-1} \frac{\sqrt{1 - \cos 2x}}{\sqrt{1 + \cos 2x}} = x$$

$$04. \tan^{-1} [\operatorname{cosec} x - \cot x] = x/2$$

$$05. \tan^{-1} \sqrt{\frac{1 - \cos x}{1 + \cos x}} = x/2$$

$$06. \tan^{-1} \left(\frac{\cos x - \sin x}{\cos x + \sin x} \right) = \pi/4 - x$$

$$07. \cot^{-1} \sqrt{\frac{1 - \sin x}{1 + \sin x}} = \pi/4 + x/2$$

$$08. \tan^{-1} \left(\frac{\cos x}{1 - \sin x} \right) = \pi/4 + x/2$$

$$09. \tan^{-1} \left(\frac{a \cos x - b \sin x}{b \cos x + a \sin x} \right) = x$$

Q SET - 4

$$01. \cos^{-1} (4x^3 - 3x) = 3\cos^{-1}x$$

$$02. \cos^{-1} \left(\frac{1 - x^2}{1 + x^2} \right) = 2 \tan^{-1} x$$

$$03. \sin^{-1} \left(\frac{1 - x^2}{1 + x^2} \right) = \frac{\pi}{2} - 2 \tan^{-1} x$$

$$04. \tan^{-1} \left(\frac{3x - x^3}{1 - 3x^2} \right) = 3 \tan^{-1} x$$

$$05. \tan^{-1} \left(\frac{x}{\sqrt{a^2 - x^2}} \right) = \sin^{-1} \left(\frac{x}{a} \right)$$

SOLUTION TO Q SET - 1

$$01. \tan^{-1}\left[\frac{2}{3}\right] + \tan^{-1}\left[\frac{1}{5}\right] = \frac{\pi}{4}$$

$$= \tan^{-1}\left(\frac{\frac{2}{3} + \frac{1}{5}}{1 - \frac{2}{3} \cdot \frac{1}{5}}\right)$$

$$= \tan^{-1}\left(\frac{\frac{10+3}{15}}{\frac{15-2}{15}}\right)$$

$$= \tan^{-1}\left[\frac{13}{13}\right]$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

$$02. \tan^{-1}\left[\frac{7}{9}\right] + \tan^{-1}\left[\frac{1}{8}\right] = \frac{\pi}{4}$$

$$= \tan^{-1}\left(\frac{\frac{7}{9} + \frac{1}{8}}{1 - \frac{7}{9} \cdot \frac{1}{8}}\right)$$

$$= \tan^{-1}\left(\frac{\frac{56+9}{72}}{\frac{72-7}{72}}\right)$$

$$= \tan^{-1}\left[\frac{65}{65}\right]$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

$$03. \tan^{-1}\left[\frac{3}{5}\right] + \tan^{-1}\left[\frac{1}{4}\right] = \frac{\pi}{4}$$

$$= \tan^{-1}\left(\frac{\frac{3}{5} + \frac{1}{4}}{1 - \frac{3}{5} \cdot \frac{1}{4}}\right)$$

$$= \tan^{-1}\left(\frac{\frac{12+5}{20}}{\frac{20-3}{20}}\right)$$

$$= \tan^{-1}\left[\frac{17}{17}\right]$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

$$04. \cot^{-1}[8] + \cot^{-1}\left[\frac{9}{7}\right] = \frac{\pi}{4}$$

$$= \tan^{-1}\left[\frac{1}{8}\right] + \tan^{-1}\left[\frac{7}{9}\right] = \frac{\pi}{4}$$

$$= \tan^{-1}\left(\frac{\frac{1}{8} + \frac{7}{9}}{1 - \frac{1}{8} \cdot \frac{7}{9}}\right)$$

$$= \tan^{-1}\left(\frac{\frac{9+56}{72}}{\frac{72-7}{72}}\right)$$

$$= \tan^{-1}\left[\frac{65}{65}\right]$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

05.

$$\tan^{-1} \left(\frac{1}{2} \right) + \tan^{-1} \left(\frac{1}{5} \right) + \tan^{-1} \left(\frac{1}{8} \right) = \frac{\pi}{4}$$

LHS

$$= \tan^{-1} \left(\frac{\frac{1}{2} + \frac{1}{5}}{1 - \frac{1}{2} \cdot \frac{1}{5}} \right) + \tan^{-1} \left(\frac{1}{8} \right)$$

$$= \tan^{-1} \left(\frac{\frac{5+2}{10}}{\frac{10-1}{10}} \right) + \tan^{-1} \left(\frac{1}{8} \right)$$

$$= \tan^{-1} \left(\frac{7}{9} \right) + \tan^{-1} \left(\frac{1}{8} \right)$$

$$= \tan^{-1} \left(\frac{\frac{7}{9} + \frac{1}{8}}{1 - \frac{7}{9} \cdot \frac{1}{8}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{56+9}{72}}{\frac{72-7}{72}} \right)$$

$$= \tan^{-1} \left(\frac{65}{65} \right)$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

06.

$$\tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{3}{5} \right) - \tan^{-1} \left(\frac{8}{19} \right) = \frac{\pi}{4}$$

LHS

$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{3}{5}}{1 - \frac{3}{4} \cdot \frac{3}{5}} \right) - \tan^{-1} \left(\frac{8}{19} \right)$$

$$= \tan^{-1} \left(\frac{\frac{15+12}{20}}{\frac{20-9}{10}} \right) - \tan^{-1} \left(\frac{8}{19} \right)$$

$$= \tan^{-1} \left(\frac{27}{11} \right) - \tan^{-1} \left(\frac{8}{19} \right)$$

$$= \tan^{-1} \left(\frac{\frac{27}{11} - \frac{8}{19}}{1 + \frac{27}{11} \cdot \frac{8}{19}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{513-88}{88}}{\frac{209+216}{88}} \right)$$

$$= \tan^{-1} \left(\frac{425}{425} \right)$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

07.

$$2\tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{4} \right) = \tan^{-1} \left(\frac{16}{13} \right)$$

LHS

$$= \tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{4} \right)$$

$$= \tan^{-1} \left(\frac{\frac{1}{3} + \frac{1}{3}}{1 - \frac{1}{3} \cdot \frac{1}{3}} \right) + \tan^{-1} \left(\frac{1}{4} \right)$$

$$= \tan^{-1} \left(\frac{\frac{3+3}{9}}{\frac{9-1}{9}} \right) + \tan^{-1} \left(\frac{1}{4} \right)$$

$$= \tan^{-1} \left(\frac{6}{8} \right) + \tan^{-1} \left(\frac{1}{4} \right)$$

$$= \tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{1}{4} \right)$$

$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{1}{4}}{1 - \frac{3}{4} \cdot \frac{1}{4}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{12+4}{12}}{\frac{16-3}{72}} \right)$$

$$= \tan^{-1} \left(\frac{16}{13} \right)$$

08.

$$2\tan^{-1} \left(\frac{1}{8} \right) + \tan^{-1} \left(\frac{1}{7} \right) + 2\tan^{-1} \left(\frac{1}{5} \right) = \frac{\pi}{4}$$

$$2\tan^{-1} \left(\frac{1}{8} \right) + 2\tan^{-1} \left(\frac{1}{5} \right) + \tan^{-1} \left(\frac{1}{7} \right) = \frac{\pi}{4}$$

LHS

$$= 2\tan^{-1} \left(\frac{\frac{1}{8} + \frac{1}{5}}{1 - \frac{1}{8} \cdot \frac{1}{5}} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= 2\tan^{-1} \left(\frac{\frac{5+8}{40}}{\frac{40-1}{40}} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= 2\tan^{-1} \left(\frac{13}{39} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= 2\tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= \tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{3} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= \tan^{-1} \left(\frac{\frac{1}{3} + \frac{1}{3}}{1 - \frac{1}{3} \cdot \frac{1}{3}} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= \tan^{-1} \left(\frac{\frac{3+3}{9}}{\frac{9-1}{9}} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= \tan^{-1} \left(\frac{6}{8} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= \tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{1}{7} \right)$$

$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{1}{7}}{1 - \frac{3}{4} \cdot \frac{1}{7}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{21+4}{28}}{\frac{28-3}{72}} \right)$$

$$= \tan^{-1} \left(\frac{25}{25} \right)$$

$$= \tan^{-1}(1) = \pi / 4$$

SOLUTION TO Q SET - 2

$$01. \cos^{-1} \left[\frac{4}{5} \right] + \cos^{-1} \left[\frac{12}{13} \right] = \cos^{-1} \left[\frac{33}{65} \right]$$

STEP 1

$$\cos^{-1} \left[\frac{4}{5} \right] = A$$

$$\frac{4}{5} = \cos A$$

$$\cos^2 A + \sin^2 A = 1$$

$$\frac{16}{25} + \sin^2 A = 1$$

$$\sin^2 A = 1 - \frac{16}{25}$$

$$\sin^2 A = \frac{9}{25}$$

$$\sin A = \frac{3}{5}$$

$$\tan A = \frac{\sin A}{\cos A}$$

$$\tan A = \frac{3}{4}$$

$$A = \tan^{-1} \left[\frac{3}{4} \right]$$

$$\text{HENCE } \cos^{-1} \left[\frac{4}{5} \right] = \tan^{-1} \left[\frac{3}{4} \right]$$

STEP 2

$$\cos^{-1} \left[\frac{12}{13} \right] = B$$

$$\frac{12}{13} = \cos B$$

$$\cos^2 B + \sin^2 B = 1$$

$$\frac{144}{169} + \sin^2 B = 1$$

$$\sin^2 B = 1 - \frac{144}{169}$$

$$\sin^2 B = \frac{25}{169}$$

$$\sin B = \frac{5}{13}$$

$$\tan B = \frac{\sin B}{\cos B}$$

$$\tan B = \frac{5}{12}$$

$$B = \tan^{-1} \left[\frac{5}{12} \right]$$

$$\text{HENCE } \cos^{-1} \left[\frac{12}{13} \right] = \tan^{-1} \left[\frac{5}{12} \right]$$

STEP 3

$$\cos^{-1} \left[\frac{33}{65} \right] = C$$

$$\frac{33}{65} = \cos C$$

$$\cos^2 C + \sin^2 C = 1$$

$$\frac{33^2}{65^2} + \sin^2 C = 1$$

$$\sin^2 C = 1 - \frac{33^2}{65^2}$$

$$\sin^2 C = \frac{65^2 - 33^2}{65^2}$$

$$\sin^2 C = \frac{(65 + 33)(65 - 33)}{65^2}$$

$$\sin^2 C = \frac{(98)(32)}{65^2}$$

$$\sin^2 C = \frac{(49)(64)}{65^2}$$

$$\sin C = \frac{7 \times 8}{65}$$

$$\sin C = \frac{56}{65}$$

$$\tan C = \frac{\sin C}{\cos C}$$

$$\tan C = \frac{56}{33}$$

$$C = \tan^{-1} \left(\frac{56}{33} \right)$$

HENCE $\cos^{-1} \left(\frac{33}{65} \right) = \tan^{-1} \left(\frac{56}{33} \right)$

STEP 4 :

$$\cos^{-1} \left(\frac{4}{5} \right) + \cos^{-1} \left(\frac{12}{13} \right) = \cos^{-1} \left(\frac{33}{65} \right)$$

WE PROVE

$$\tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{5}{12} \right) = \tan^{-1} \left(\frac{56}{33} \right)$$

LHS

$$= \tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{5}{12} \right)$$

$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{5}{12}}{1 - \frac{3}{4} \cdot \frac{5}{12}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{36+20}{84}}{\frac{48-15}{84}} \right)$$

$$= \tan^{-1} \left(\frac{56}{33} \right)$$

= RHS

02. $\sin^{-1} \left(\frac{3}{5} \right) + \cos^{-1} \left(\frac{12}{13} \right) = \sin^{-1} \left(\frac{56}{65} \right)$

STEP 1

$$\sin^{-1} \left(\frac{3}{5} \right) = A$$

$$\frac{3}{5} = \sin A$$

$$\sin^2 A + \cos^2 A = 1$$

$$\frac{9}{25} + \cos^2 A = 1$$

$$\cos^2 A = 1 - \frac{9}{25}$$

$$\cos^2 A = \frac{16}{25}$$

$$\cos A = \frac{4}{5}$$

$$\tan A = \frac{\sin A}{\cos A}$$

$$\tan A = \frac{3}{4}$$

$$A = \tan^{-1} \left(\frac{3}{4} \right)$$

HENCE $\sin^{-1} \left(\frac{3}{5} \right) = \tan^{-1} \left(\frac{3}{4} \right)$

STEP 2

$$\cos^{-1} \left(\frac{12}{13} \right) = B$$

$$\frac{12}{13} = \cos B$$

$$\cos^2 B + \sin^2 B = 1$$

$$\frac{144}{169} + \sin^2 B = 1$$

$$\sin^2 B = 1 - \frac{144}{169}$$

$$\sin^2 B = \frac{25}{169}$$

$$\sin B = \frac{5}{13}$$

$$\tan B = \frac{\sin B}{\cos B}$$

$$\tan B = \frac{5}{12}$$

$$B = \tan^{-1} \left(\frac{5}{12} \right)$$

$$\text{HENCE } \cos^{-1} \left(\frac{12}{13} \right) = \tan^{-1} \left(\frac{5}{12} \right)$$

STEP 3

$$\sin^{-1} \left(\frac{56}{65} \right) = C$$

$$\frac{56}{65} = \sin C$$

$$\sin^2 C + \cos^2 C = 1$$

$$\frac{56^2}{65^2} + \cos^2 C = 1$$

$$\cos^2 C = 1 - \frac{56^2}{65^2}$$

$$\cos^2 C = \frac{65^2 - 56^2}{65^2}$$

$$\cos^2 C = \frac{(65 + 56)(65 - 56)}{65^2}$$

$$\cos^2 C = \frac{(121)(9)}{65^2}$$

$$\cos C = \frac{11 \times 3}{65}$$

$$\cos C = \frac{33}{65}$$

$$\tan C = \frac{\sin C}{\cos C}$$

$$\tan C = \frac{56}{33}$$

$$C = \tan^{-1} \left(\frac{56}{33} \right)$$

$$\text{HENCE } \sin^{-1} \left(\frac{56}{65} \right) = \tan^{-1} \left(\frac{56}{33} \right)$$

STEP 4 :

$$\sin^{-1} \left(\frac{3}{5} \right) + \cos^{-1} \left(\frac{12}{13} \right) = \sin^{-1} \left(\frac{56}{65} \right)$$

WE PROVE

$$\tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{5}{12} \right) = \tan^{-1} \left(\frac{56}{33} \right)$$

LHS

$$= \tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{5}{12} \right)$$

$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{5}{12}}{1 - \frac{3}{4} \cdot \frac{5}{12}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{36+20}{84}}{\frac{48-15}{84}} \right)$$

$$= \tan^{-1} \left(\frac{56}{33} \right)$$

$$= \text{RHS}$$

$$03. \quad \sin^{-1} \left[\frac{3}{5} \right] + \sin^{-1} \left[\frac{8}{17} \right] = \sin^{-1} \left[\frac{77}{85} \right]$$

STEP 1

$$\sin^{-1} \left[\frac{3}{5} \right] = A$$

$$\frac{3}{5} = \sin A$$

$$\sin^2 A + \cos^2 A = 1$$

$$\frac{9}{25} + \cos^2 A = 1$$

$$\cos^2 A = 1 - \frac{9}{25}$$

$$\cos^2 A = \frac{16}{25}$$

$$\cos A = \frac{4}{5}$$

$$\tan A = \frac{\sin A}{\cos A}$$

$$\tan A = \frac{3}{4}$$

$$A = \tan^{-1} \left[\frac{3}{4} \right]$$

$$\text{HENCE } \sin^{-1} \left[\frac{3}{5} \right] = \tan^{-1} \left[\frac{3}{4} \right]$$

STEP 2

$$\sin^{-1} \left[\frac{8}{17} \right] = B$$

$$\frac{8}{17} = \sin B$$

$$\sin^2 B + \cos^2 B = 1$$

$$\frac{64}{289} + \cos^2 B = 1$$

$$\cos^2 B = 1 - \frac{64}{289}$$

$$\cos^2 B = \frac{225}{289}$$

$$\cos B = \frac{15}{17}$$

$$\tan B = \frac{\sin B}{\cos B}$$

$$\tan B = \frac{8}{15}$$

$$B = \tan^{-1} \left[\frac{8}{15} \right]$$

$$\text{HENCE } \sin^{-1} \left[\frac{8}{17} \right] = \tan^{-1} \left[\frac{8}{15} \right]$$

STEP 3

$$\sin^{-1} \left[\frac{77}{85} \right] = C$$

$$\frac{77}{85} = \sin C$$

$$\sin^2 C + \cos^2 C = 1$$

$$\frac{77^2}{85^2} + \cos^2 C = 1$$

$$\cos^2 C = 1 - \frac{77^2}{85^2}$$

$$\cos^2 C = \frac{85^2 - 77^2}{85^2}$$

$$\cos^2 C = \frac{(85 + 77)(85 - 77)}{85^2}$$

$$\cos^2 C = \frac{(162)(8)}{85^2}$$

$$\cos^2 C = \frac{(81)(16)}{85^2}$$

$$\cos C = \frac{9 \times 4}{85}$$

$$\cos C = \frac{36}{85}$$

$$\tan C = \frac{\sin C}{\cos C}$$

$$\tan C = \frac{77}{36}$$

$$C = \tan^{-1} \left(\frac{77}{36} \right)$$

HENCE $\sin^{-1} \left(\frac{77}{85} \right) = \tan^{-1} \left(\frac{77}{36} \right)$

STEP 4 :

$$\sin^{-1} \left(\frac{3}{5} \right) + \sin^{-1} \left(\frac{8}{17} \right) = \sin^{-1} \left(\frac{77}{85} \right)$$

WE PROVE

$$\tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{8}{15} \right) = \tan^{-1} \left(\frac{77}{36} \right)$$

LHS

$$= \tan^{-1} \left(\frac{3}{4} \right) + \tan^{-1} \left(\frac{8}{15} \right)$$

$$= \tan^{-1} \left(\frac{\frac{3}{4} + \frac{8}{15}}{1 - \frac{3}{4} \cdot \frac{8}{15}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{45+32}{60}}{\frac{60-24}{60}} \right)$$

$$= \tan^{-1} \left(\frac{77}{36} \right)$$

= RHS

$$04. \quad \cos^{-1} \left(\frac{12}{13} \right) + \sin^{-1} \left(\frac{5}{13} \right) = \cot^{-1} \left(\frac{119}{120} \right)$$

STEP 1

$$\cos^{-1} \left(\frac{12}{13} \right) = A$$

$$\frac{12}{13} = \cos A$$

$$\cos^2 A + \sin^2 A = 1$$

$$\frac{144}{169} + \sin^2 A = 1$$

$$\sin^2 A = 1 - \frac{144}{169}$$

$$\sin^2 A = \frac{25}{169}$$

$$\sin A = \frac{5}{13}$$

$$\tan A = \frac{\sin A}{\cos A}$$

$$\tan A = \frac{5}{12}$$

$$A = \tan^{-1} \left(\frac{5}{12} \right)$$

$$\text{HENCE } \cos^{-1} \left(\frac{12}{13} \right) = \tan^{-1} \left(\frac{5}{12} \right)$$

STEP 2

$$\sin^{-1} \left(\frac{5}{13} \right) = B$$

$$\frac{5}{13} = \sin B$$

$$\sin^2 B + \cos^2 B = 1$$

$$\frac{25}{169} + \cos^2 B = 1$$

$$\cos^2 B = 1 - \frac{25}{169}$$

$$\cos^2 B = \frac{144}{169}$$

$$\cos B = \frac{12}{13}$$

$$\tan B = \frac{\sin B}{\cos B}$$

$$\tan B = \frac{5}{12}$$

$$B = \tan^{-1} \left(\frac{5}{12} \right)$$

$$\text{HENCE } \sin^{-1} \left(\frac{5}{13} \right) = \tan^{-1} \left(\frac{5}{12} \right)$$

STEP 3 :

$$\cos^{-1} \left(\frac{12}{13} \right) + \sin^{-1} \left(\frac{5}{13} \right) = \cot^{-1} \left(\frac{119}{120} \right)$$

WE PROVE

$$\tan^{-1} \left(\frac{5}{12} \right) + \tan^{-1} \left(\frac{5}{12} \right) = \tan^{-1} \left(\frac{120}{119} \right)$$

LHS

$$= \tan^{-1} \left(\frac{5}{12} \right) + \tan^{-1} \left(\frac{5}{12} \right)$$

$$= \tan^{-1} \left(\frac{\frac{5}{12} + \frac{5}{12}}{1 - \frac{5}{12} \cdot \frac{5}{12}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{60+60}{144}}{\frac{144-25}{144}} \right)$$

$$= \tan^{-1} \left(\frac{120}{119} \right)$$

= RHS

05.

$$2\sin^{-1}\left[\frac{3}{5}\right] - \tan^{-1}\left[\frac{17}{31}\right] = \frac{\pi}{4}$$

STEP 1

$$\sin^{-1}\left[\frac{3}{5}\right] = A$$

$$\frac{3}{5} = \sin A$$

$$\sin^2 A + \cos^2 A = 1$$

$$\frac{9}{25} + \cos^2 A = 1$$

$$\cos^2 A = 1 - \frac{9}{25}$$

$$\cos^2 A = \frac{16}{25}$$

$$\cos A = \frac{4}{5}$$

$$\tan A = \frac{\sin A}{\cos A}$$

$$\tan A = \frac{3}{4}$$

$$A = \tan^{-1}\left[\frac{3}{4}\right]$$

HENCE $\sin^{-1}\left[\frac{3}{5}\right] = \tan^{-1}\left[\frac{3}{4}\right]$

STEP 2 :

$$2\sin^{-1}\left[\frac{3}{5}\right] - \tan^{-1}\left[\frac{17}{31}\right] = \frac{\pi}{4}$$

We Prove

$$2\tan^{-1}\left[\frac{3}{4}\right] - \tan^{-1}\left[\frac{17}{31}\right] = \frac{\pi}{4}$$

LHS

$$2\tan^{-1}\left[\frac{3}{4}\right] - \tan^{-1}\left[\frac{17}{31}\right]$$

$$= \tan^{-1}\left[\frac{3}{4}\right] + \tan^{-1}\left[\frac{3}{4}\right] - \tan^{-1}\left[\frac{17}{31}\right]$$

$$= \tan^{-1}\left[\frac{\frac{3}{4} + \frac{3}{4}}{1 - \frac{3}{4} \cdot \frac{3}{4}}\right] - \tan^{-1}\left[\frac{17}{31}\right]$$

$$= \tan^{-1}\left[\frac{\frac{12+12}{16}}{\frac{16-9}{16}}\right] - \tan^{-1}\left[\frac{17}{31}\right]$$

$$= \tan^{-1}\left[\frac{24}{7}\right] - \tan^{-1}\left[\frac{17}{31}\right]$$

$$= \tan^{-1}\left[\frac{\frac{24}{7} - \frac{17}{31}}{1 + \frac{24}{7} \cdot \frac{17}{31}}\right]$$

$$= \tan^{-1}\left[\frac{\frac{744-119}{217}}{\frac{217+408}{217}}\right]$$

$$= \tan^{-1}\left[\frac{625}{625}\right]$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

$$06. \quad \sin^{-1} \left(\frac{3}{\sqrt{34}} \right) + \cos^{-1} \left(\frac{4}{\sqrt{17}} \right) = \frac{\pi}{4}$$

STEP 1

$$\sin^{-1} \left(\frac{3}{\sqrt{34}} \right) = A$$

$$\frac{3}{\sqrt{34}} = \sin A$$

$$\sin^2 A + \cos^2 A = 1$$

$$\frac{9}{34} + \cos^2 A = 1$$

$$\cos^2 A = 1 - \frac{9}{34}$$

$$\cos^2 A = \frac{25}{34}$$

$$\cos A = \frac{5}{\sqrt{34}}$$

$$\tan A = \frac{\sin A}{\cos A}$$

$$\tan A = \frac{3}{5}$$

$$A = \tan^{-1} \left(\frac{3}{5} \right)$$

$$\text{HENCE } \sin^{-1} \left(\frac{3}{\sqrt{34}} \right) = \tan^{-1} \left(\frac{3}{5} \right)$$

STEP 2

$$\cos^{-1} \left(\frac{4}{\sqrt{17}} \right) = B$$

$$\frac{4}{\sqrt{17}} = \cos B$$

$$\cos^2 B + \sin^2 B = 1$$

$$\frac{16}{17} + \sin^2 B = 1$$

$$\sin^2 B = 1 - \frac{16}{17}$$

$$\sin^2 B = \frac{1}{17}$$

$$\sin B = \frac{1}{\sqrt{17}}$$

$$\tan B = \frac{\sin B}{\cos B}$$

$$\tan B = \frac{1}{4}$$

$$B = \tan^{-1} \left(\frac{1}{4} \right)$$

$$\text{HENCE } \cos^{-1} \left(\frac{4}{\sqrt{17}} \right) = \tan^{-1} \left(\frac{1}{4} \right)$$

STEP 3

$$\sin^{-1} \left(\frac{3}{\sqrt{34}} \right) + \cos^{-1} \left(\frac{4}{\sqrt{17}} \right) = \frac{\pi}{4}$$

We Prove

$$\tan^{-1} \left(\frac{3}{5} \right) + \tan^{-1} \left(\frac{1}{4} \right) = \frac{\pi}{4}$$

$$= \tan^{-1} \left(\frac{\frac{3}{5} + \frac{1}{4}}{1 - \frac{3}{5} \cdot \frac{1}{4}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{12+5}{20}}{\frac{20-3}{20}} \right)$$

$$= \tan^{-1} \left(\frac{17}{17} \right)$$

$$= \tan^{-1}(1)$$

$$= \pi/4$$

$$07. \quad \sin^{-1} \left(\frac{2}{\sqrt{13}} \right) + \cos^{-1} \left(\frac{5}{\sqrt{26}} \right) = \frac{\pi}{4}$$

STEP 1

$$\sin^{-1} \left(\frac{2}{\sqrt{13}} \right) = A$$

$$\frac{2}{\sqrt{13}} = \sin A$$

$$\sin^2 A + \cos^2 A = 1$$

$$\frac{4}{13} + \cos^2 A = 1$$

$$\cos^2 A = 1 - \frac{4}{13}$$

$$\cos^2 A = \frac{9}{13}$$

$$\cos A = \frac{3}{\sqrt{13}}$$

$$\tan A = \frac{\sin A}{\cos A}$$

$$\tan A = \frac{2}{3}$$

$$A = \tan^{-1} \left(\frac{2}{3} \right)$$

$$\text{HENCE } \sin^{-1} \left(\frac{2}{\sqrt{13}} \right) = \tan^{-1} \left(\frac{2}{3} \right)$$

STEP 2

$$\cos^{-1} \left(\frac{5}{\sqrt{26}} \right) = B$$

$$\frac{5}{\sqrt{26}} = \cos B$$

$$\cos^2 B + \sin^2 B = 1$$

$$\frac{25}{26} + \sin^2 B = 1$$

$$\sin^2 B = 1 - \frac{25}{26}$$

$$\sin^2 B = \frac{1}{26}$$

$$\sin B = \frac{1}{\sqrt{26}}$$

$$\tan B = \frac{\sin B}{\cos B}$$

$$\tan B = \frac{1}{5}$$

$$B = \tan^{-1} \left(\frac{1}{5} \right)$$

$$\text{HENCE } \cos^{-1} \left(\frac{5}{\sqrt{26}} \right) = \tan^{-1} \left(\frac{1}{5} \right)$$

STEP 3

$$\sin^{-1} \left(\frac{2}{\sqrt{13}} \right) + \cos^{-1} \left(\frac{5}{\sqrt{26}} \right) = \frac{\pi}{4}$$

We Prove

$$\tan^{-1} \left(\frac{2}{3} \right) + \tan^{-1} \left(\frac{1}{5} \right) = \frac{\pi}{4}$$

$$= \tan^{-1} \left(\frac{\frac{2}{3} + \frac{1}{5}}{1 - \frac{2}{3} \cdot \frac{1}{5}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{10+3}{15}}{\frac{15-2}{15}} \right)$$

$$= \tan^{-1} \left(\frac{13}{13} \right)$$

$$= \tan^{-1} (1)$$

$$= \pi/4$$

SOLUTION TO Q SET - 3

$$\begin{aligned}
 01. \tan^{-1} \left(\frac{\sin 2x}{1 + \cos 2x} \right) &= x \\
 &= \tan^{-1} \left(\frac{2 \sin x \cdot \cos x}{2 \cos^2 x} \right) \\
 &= \tan^{-1} \left(\frac{\sin x}{\cos x} \right) \\
 &= \tan^{-1} [\tan x] \\
 &= x \qquad \qquad \qquad = \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 02. \cot^{-1} \left(\frac{\sin 2x}{1 - \cos 2x} \right) &= x \\
 &= \cot^{-1} \left(\frac{2 \sin x \cdot \cos x}{2 \sin^2 x} \right) \\
 &= \cot^{-1} \left(\frac{\cos x}{\sin x} \right) \\
 &= \cot^{-1} [\cot x] \\
 &= x \qquad \qquad \qquad = \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 03. \tan^{-1} \sqrt{\frac{1 - \cos 2x}{1 + \cos 2x}} &= x \\
 &= \tan^{-1} \sqrt{\frac{2 \sin^2 x}{2 \cos^2 x}} \\
 &= \tan^{-1} \frac{\sin x}{\cos x} \\
 &= \tan^{-1} [\tan x] \\
 &= x \qquad \qquad \qquad = \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 04. \tan^{-1} [\operatorname{cosec} x - \cot x] &= x/2 \\
 &= \tan^{-1} \left(\frac{1 - \cos x}{\sin x} \right) \\
 &= \tan^{-1} \left(\frac{2 \sin^2 x/2}{2 \sin x/2 \cdot \cos x/2} \right) \\
 &= \tan^{-1} \left(\frac{\sin x/2}{\cos x/2} \right) \\
 &= \tan^{-1} [\tan x/2] \\
 &= x/2 \qquad \qquad \qquad = \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 05. \tan^{-1} \sqrt{\frac{1 - \cos x}{1 + \cos x}} &= x/2 \\
 &= \tan^{-1} \sqrt{\frac{2 \sin^2 x/2}{2 \cos^2 x/2}} \\
 &= \tan^{-1} \left(\frac{\sin x/2}{\cos x/2} \right) \\
 &= \tan^{-1} [\tan x/2] \\
 &= x/2 \qquad \qquad \qquad = \text{RHS}
 \end{aligned}$$

$$\begin{aligned}
 06. \tan^{-1} \left(\frac{\cos x - \sin x}{\cos x + \sin x} \right) &= \pi/4 - x \\
 &= \tan^{-1} \left(\frac{\frac{\cos x - \sin x}{\cos x}}{\frac{\cos x + \sin x}{\cos x}} \right)
 \end{aligned}$$

$$= \tan^{-1} \left(\frac{1 - \tan x}{1 + \tan x} \right)$$

$$= \tan^{-1} \left(\frac{\tan \pi/4 - \tan x}{1 + \tan \pi/4 \cdot \tan x} \right)$$

$$= \tan^{-1} \tan (\pi/4 - x)$$

$$= (\pi/4 - x) = \text{RHS}$$

$$07. \quad \cot^{-1} \sqrt{\frac{1 - \sin x}{1 + \sin x}} = \pi/4 + x/2$$

$$= \tan^{-1} \sqrt{\frac{1 + \sin x}{1 - \sin x}}$$

$$= \tan^{-1} \sqrt{\frac{\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2} + 2 \cdot \sin \frac{x}{2} \cdot \cos \frac{x}{2}}{\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2} - 2 \cdot \sin \frac{x}{2} \cdot \cos \frac{x}{2}}}$$

$$= \tan^{-1} \sqrt{\frac{(\cos x/2 + \sin x/2)^2}{(\cos x/2 - \sin x/2)^2}}$$

$$= \tan^{-1} \left(\frac{\cos x/2 + \sin x/2}{\cos x/2 - \sin x/2} \right)$$

$$= \tan^{-1} \left(\frac{\frac{\cos x/2 + \sin x/2}{\cos x/2}}{\frac{\cos x/2 - \sin x/2}{\cos x/2}} \right)$$

$$= \tan^{-1} \left(\frac{1 + \tan x/2}{1 - \tan x/2} \right)$$

$$= \tan^{-1} \left(\frac{\tan \pi/4 + \tan x/2}{1 - \tan \pi/4 \cdot \tan x/2} \right)$$

$$= \tan^{-1} \tan (\pi/4 + x/2)$$

$$= \pi/4 + x/2 = \text{RHS}$$

$$08. \quad \tan^{-1} \left(\frac{\cos x}{1 - \sin x} \right) = \pi/4 + x/2$$

$$= \tan^{-1} \left(\frac{\frac{\cos^2 \frac{x}{2} - \sin^2 \frac{x}{2}}{2}}{\frac{\cos^2 \frac{x}{2} + \sin^2 \frac{x}{2} - 2 \cdot \sin \frac{x}{2} \cdot \cos \frac{x}{2}}{2}} \right)$$

$$= \tan^{-1} \left(\frac{(\cos x/2 - \sin x/2)(\cos x/2 + \sin x/2)}{(\cos x/2 - \sin x/2)^2} \right)$$

$$= \tan^{-1} \left(\frac{\cos x/2 + \sin x/2}{\cos x/2 - \sin x/2} \right)$$

$$= \tan^{-1} \left(\frac{\frac{\cos x/2 + \sin x/2}{\cos x/2}}{\frac{\cos x/2 - \sin x/2}{\cos x/2}} \right)$$

$$= \tan^{-1} \left(\frac{1 + \tan x/2}{1 - \tan x/2} \right)$$

$$= \tan^{-1} \left(\frac{\tan \pi/4 + \tan x/2}{1 - \tan \pi/4 \cdot \tan x/2} \right)$$

$$= \tan^{-1} \tan (\pi/4 + x/2)$$

$$= \pi/4 + x/2 = \text{RHS}$$

$$09. \quad \tan^{-1} \left(\frac{a \cos x - b \sin x}{b \cos x + a \sin x} \right) = x$$

$$= \tan^{-1} \left(\frac{\frac{a \cos x - b \sin x}{b \cos x}}{\frac{b \cos x + a \sin x}{b \cos x}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{a \cos x}{b \cos x} - \frac{b \sin x}{b \cos x}}{\frac{b \cos x}{b \cos x} + \frac{a \sin x}{b \cos x}} \right)$$

$$= \tan^{-1} \left(\frac{\frac{a}{b} - \tan x}{1 + \frac{a}{b} \cdot \tan x} \right)$$

$$= \tan^{-1} \frac{a}{b} - \tan^{-1} \tan x$$

$$= \tan^{-1} \frac{a}{b} - x$$

SOLUTION TO Q SET - 4

$$01. \quad \cos^{-1} (4x^3 - 3x) = 3\cos^{-1}x$$

LHS

$$= \cos^{-1} (4x^3 - 3x)$$

$$\text{Put } x = \cos \theta$$

$$= \cos^{-1} (4\cos^3 \theta - 3\cos \theta)$$

$$= \cos^{-1} (\cos 3\theta)$$

$$= 3\theta$$

$$= 3 \cos^{-1}x = \text{RHS}$$

$$02. \quad \cos^{-1} \left(\frac{1 - x^2}{1 + x^2} \right) = 2 \tan^{-1} x$$

LHS

$$= \cos^{-1} \left(\frac{1 - x^2}{1 + x^2} \right)$$

$$\text{Put } x = \tan \theta$$

$$= \cos^{-1} \left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right)$$

$$= \cos^{-1} \cos 2\theta$$

$$= 2\theta$$

$$= 2 \tan^{-1} x = \text{RHS}$$

$$03. \quad \sin^{-1} \left(\frac{1 - x^2}{1 + x^2} \right) = \frac{\pi}{2} - 2 \tan^{-1} x$$

LHS

$$= \sin^{-1} \left(\frac{1 - x^2}{1 + x^2} \right)$$

$$\text{Put } x = \tan \theta$$

$$= \cos^{-1} \left(\frac{1 - \tan^2 \theta}{1 + \tan^2 \theta} \right)$$

$$= \sin^{-1} \cos 2\theta$$

$$= \sin^{-1} \sin (\pi/2 - 2\theta)$$

$$= \pi/2 - 2\theta$$

$$= \pi/2 - 2 \tan^{-1} x = \text{RHS}$$

$$04. \quad \tan^{-1} \left(\frac{3x - x^3}{1 + 3x^2} \right) = 3 \tan^{-1} x$$

LHS

$$= \tan^{-1} \left(\frac{3x - x^3}{1 + 3x^2} \right)$$

Put $x = \tan \theta$

$$= \tan^{-1} \left(\frac{3 \tan \theta - \tan^3 \theta}{1 + 3 \tan^2 \theta} \right)$$

$$= \tan^{-1} \tan 3\theta$$

$$= 3\theta$$

$$= 3 \tan^{-1} x = \text{RHS}$$

05. $\tan^{-1} \left(\frac{x}{\sqrt{a^2 - x^2}} \right) = \sin^{-1} \left(\frac{x}{a} \right)$

LHS

$$= \tan^{-1} \left(\frac{x}{\sqrt{a^2 - x^2}} \right)$$

Put $x = a \sin \theta$

$$= \tan^{-1} \left(\frac{a \sin \theta}{\sqrt{a^2 - a^2 \sin^2 \theta}} \right)$$

$$= \tan^{-1} \left(\frac{a \sin \theta}{\sqrt{a^2 (1 - \sin^2 \theta)}} \right)$$

$$= \tan^{-1} \left(\frac{a \sin \theta}{\sqrt{a^2 \cos^2 \theta}} \right)$$

$$= \tan^{-1} \left(\frac{a \sin \theta}{a \cos \theta} \right)$$

$$= \tan^{-1} \tan \theta$$

$$= \theta \longrightarrow$$

$$= \sin^{-1} \left(\frac{x}{a} \right) \longleftarrow$$

$$= \text{RHS}$$

EXTRA

$$\tan^{-1} 1 + \tan^{-1} 2 + \tan^{-1} 3 = \pi$$

NOTE

if $xy > 1$ THEN

$$\tan^{-1} x + \tan^{-1} y = \pi + \tan^{-1} \left(\frac{x+y}{1-xy} \right)$$

LHS

$$= \tan^{-1} 1 + \pi + \tan^{-1} \left(\frac{2+3}{1-2.3} \right)$$

$$= \tan^{-1} 1 + \pi + \tan^{-1} \left(\frac{5}{-5} \right)$$

$$= \tan^{-1} 1 + \pi + \tan^{-1} (-1)$$

$$= \tan^{-1} 1 + \pi - \tan^{-1} 1$$

$$= \pi$$

$x = a \cdot \sin \theta$
$\sin \theta = \frac{x}{a}$
$\theta = \sin^{-1} \frac{x}{a}$